

Sample no.	Rex 24C						
	Abundance	Distribution & Remarks	Size (voids) or thickness (laminae) in mm	Mineralogy	Abundance	Size (mm)	Shape
<b>Matrix</b>	46%	Fine grained and very finely intersperced.		Braunite	xxxx	0.004	Anhedral
				Microspar [Kutnahorite]	xxx	0.005-0.01	Anhedral
				Hematite	xx	0.001-0.002	Anhedral
				Hausmannite	xx	0.015	Euhedral to subhedral
<b>Ovoids</b>	40%	Unevenly; Accretions of ellipsoidal and rounded ovoids of small to medium size in cloud structures give the zone a banded (2mm) appearance. Ovoids consist mainly of microspar and have braunite, hematite, hausmannite and todorokite inclusions. Todorokite have hematite inclusions.	0.55x0.27 0.5x0.15	Braunite	x	0.003	Anhedral
				Pseudospar [Calcite]	xxxx	0.02	Anhedral
				Hematite	x	0.001	Anhedral
				Hausmannite	x	0.006	Euhedral
				Todorokite	x	0.009	Euhedral
<b>Laminae</b>	10%	Unevenly; Finely laminated and consists mainly of pseudospar with braunite, hematite and hausmannite inclusions. Hematite inclusions occur in the braunite and are finely intergrown with the braunite.	0.25	Braunite	x	0.0025	Anhedral
				Pseudospar [Kutnahorite]	xxxx	0.01	Anhedral
				Hematite	x	0.001	Anhedral
				Hausmannite	xx	0.0025	Anhedral
<b>Other</b>		The zone has a banded appearance					
	1%	Cross-cutting veins	0.02-0.3	Todorokite	xxxx	0.001	Needles
	3%	Porosity	0.025				

xxxx - Dominant (>50%); xxx - Major (20-50%); xx - Minor (5-20%); x - Trace (<5%)