
REFERENCES

Altermann, W., and Siegfried, H.P., (1997): Sedimentology and facies development of an Archean shelf: carbonate platform transition in the Kaapvaal Craton, as deduced from a deep borehole at Kathu, South Africa. *Journal of African Earth Sciences* 24 pp. 391-410.

Altermann, W., and Nelson, D.R., (1998): Sedimentation rates, basin analysis and regional correlations of three Neoproterozoic and Palaeoproterozoic sub-basins of the Kaapvaal craton as inferred from precise U-Pb zircon ages from volcanoclastic sediments. *Sedimentary Geology*, 120 pp. 225-256.

Andrade, F.R.D., Möller, P., Lüders, V., Dulski, P., Gilg, H.A., (1999): Hydrothermal rare earth elements mineralization in the Barra do Itapirapuã carbonatite, southern Brazil: behaviour of selected trace elements and stable isotopes (C,O). *Chemical Geology*, 155, pp. 91-113.

Barton, E.S., Altermann, W., Williams, I.S., Smith, C.B., (1994): U-Pb age for a tuff in the Campbellrand Group, Griqualand West sequence, South Africa: implications for early Proterozoic rock accumulation rates. *Geology*, 22, pp. 343-346.

Bau, M., and Dulski, P., (1996): Distribution of yttrium and rare-earth elements in the Penge and Kuruman iron-formations, Transvaal Supergroup, South Africa. *Precambrian Research*, 79, pp. 37-55

Bau, M., Romer, R.L., Lüders, V., Beukes, N.J., (1999): Pb, O, and C isotopes in silicified Mooidraai dolomite (Transvaal Supergroup, South Africa): implications for the composition of Paleoproterozoic seawater and 'dating' the increase of oxygen in the Precambrian atmosphere. *Earth Planetary Science Letters*, 174, pp. 43-57.

Beukes, N.J., (1978): *Die Karbonaatgesteentes en Ysterformasies van die Ghaap-groep van die Transvaal Supergroep in Noord-kaapland*. Unpublished. Ph.D. thesis, Rand Afrikaans University, Johannesburg, 580 pp

Beukes, N.J., (1979): Litostratigrafiese onderverdeling van die Schmidtsdrif-Subgroep van die Ghaap-Groep in Noord-Kaapland. *Transvaal Geological Society of South Africa*, 82, pp. 313-327.

Beukes, N.J., (1980): Stratigrafie en lithofasies van die Campbellrand-Subgroep van die Proterofitiese Ghaap-Groep, Noord-Kaapland. *Transvaal Geological Society of South Africa*. 83, pp 141-170.

Beukes, N.J., (1983): Paleoenvironmental settings of iron-formation in the depositional basin of the Transvaal Supergroup, South Africa. In: Trendall, A.F., and Morris, R.C., *Iron-Formations: Facts and Problems*. Amsterdam, Elsevier, pp. 131-209.

Beukes, N.J., (1984): Sedimentology of the Kuruman and Griquatown Iron-Formations, Transvaal Supergroup, Griqualand West, South Africa. *Precambrian Research*, 24, pp. 47-84.

Beukes, N.J., (1986): The Transvaal Sequence in Griqualand West. In: Anhaeusser, C.R., Maske, S., (Eds.), *Mineral Deposits of Southern Africa, Vol. I. Geological Society of South Africa*, Johannesburg, pp. 819-828.

Beukes, N.J., (1987): Facies relations, depositional environments and diagenesis in a major early Proterozoic stromatolitic carbonate platform to basinal sequence, Campbellrand Subgroup, Transvaal Supergroup, South Africa. *Sedimentary Geology*, 54, pp. 1-46.

Beukes, N.J., (1987): New evidence for thrust faulting in Griqualand West, South Africa: implications for stratigraphy and the age of red beds. *Suid Afrikaanse Tydskrif fir Geologie*, 90, pp. 378-394.

Beukes, N.J., and Klein, C., (1990): Geochemical and sedimentology of a facies transition – from microbanded to granular iron-formation – in the early Proterozoic Transvaal Supergroup, South Africa. *Precambrian Research*, 47, pp. 99-139.

Beukes, N.J., Gutzmer, J., Klein, C., (2000): *Genesis of world-class, high-grade iron and manganese deposits - what we do and don't know*. GSA Abstracts with Programs, A-3.

Beukes, N.J., Gutzmer, J., and Mukhopadhyay, J., (2002): *The geology and genesis of high-grade hematite iron ore deposits*. Iron Ore Conference, Perth. Pp. 23-29.

Button, A., (1973): The depositional history of the Wolkberg proto-basin, Transvaal. *Transvaal Geological Society of South Africa* 76, pp. 15-25.

Coetzee, L.L., (2001): *Genetic stratigraphy of the Paleoproterozoic Pretoria group in the western Transvaal*. Unpublished. M.Sc. thesis, Rand Afrikaans University, Johannesburg, 212 pp.

Coplen, T.B., Kendall, C., and Hopple, J., (1983): Comparison of stable isotope reference samples. *Nature*, 302, pp. 236-238.

Cornell, D.H., Schütte, S.S., Eglington, B.L., (1996): The Ongeluk Basaltic Andesite Formation in Griqualand West, South Africa: submarine alteration in a 2222 Ma Proterozoic sea. *Precambrian research* 79, pp. 101-124.

Cornell, D.H., Armstrong, R.A., Walraven, F., (1998): Geochronology of the Proterozoic Hartley Basalt Formation, South Africa: constraints on the Kheis tectogenesis and the Kaapvaal craton's earliest Wilson cycle. *Journal of African Earth Sciences* 26, pp 5-27.

Craig, J., and Vaughan, D.J., (1994): *Ore microscopy & ore petrography*. Second edition. John Wiley & Sons, Inc., Canada.

Dorland, H.C., (1999): *Paleoproterozoic laterite, red beds and ironstone of the Pretoria Group with reference to the history of atmospheric oxygen*. Unpublished. M.Sc. thesis, Rand Afrikaans University, Johannesburg. 147pp.

Dreyer, C.J.B., (1974): *The Nauga Haematite Ore Deposit*. Unpublished Rand Mines Limited Report. 11pp.

Gutzmer, J., and Beukes, N.J., (1998): *High grade manganese ores in the Kalahari manganese field: Characterisation and dating of ore forming events*. Unpublished. Report, Rand Afrikaans University, Johannesburg, 221 pp.

Gutzmer, J., and Beukes, N.J., (2002): *Origin of manganese and iron ores at Rooinekke, Northern Cape Province*. Unpublished. Final Project Report, Rand Afrikaans University, Johannesburg, 73 pp.

Harmer, R.E., (1992): *The geochemistry of Spitskop and related alkaline Complexes*. Ph.D. dissertation 332pp. University Cape Town, South Africa.

Hoernle, K., Tilton, G., Le Bas, M.J., Duggen, S., Garbe-Schönberg, D., (2002): Geochemistry of oceanic carbonatites compared with continental carbonatite: mantle recycling of oceanic crustal carbonate. *Contribution to Mineral Petrology*, 142, pp. 520-542.

Horstmann, U.E., and Hälbich, I.W., (1995): Chemical composition of banded iron-formations of the Griqualand West Sequence, Northern Cape Province, South Africa, in comparison with other Precambrian iron formations. *Precambrian Research*, 72, pp. 109-145.

Horstmann, U.E., and Verwoerd, W.J., (1997): Carbon and oxygen isotope variations in southern African carbonatites. *Journal of African Earth Sciences*, 25, pp. 115-136.

Horstmann, U.E., (2003): *C/O stable isotope analyses of 8 carbonate samples (dolomites) of possible carbonatite origin*. Confidential report. Stable Isotope Laboratory Project: 6820-5162, Job:03/089. Council for Geoscience, Pretoria, 6 pp.

Huang, Y.-M., Hawkesworth, C.J., van Calsteren, P., McDermott, F., (1995): Geochemical characteristics and origin of the Jacupiranga carbonatites, Brazil. *Chemical Geology*, 119, pp. 79-99.

Ionov, D., Harmer, R.E., (2002): Trace element distribution in calcite-dolomite carbonatites from Spitskop: inferences for differentiation of carbonatite magmas and the origin of carbonatites in mantle xenoliths. *Earth and Planetary Science Letters*, 198, pp. 495-510.

Kaufman, A.J., (1996): Geochemical and mineralogic effects of contact metamorphism on banded iron-formation: an example from the Transvaal Basin, South Africa. *Precambrian Research* 79, pp. 171-194.

Keller, J., and Hoefs, J., (1995): Stable isotope characteristics of recent natrocarbonatites from Oldoinyo Lengai. In: Bell, K. and Keller, J., (Eds), *Carbonatite Volcanism: Oldoinyo Lengai and the petrogenesis of natrocarbonatites*. Springer Verlag, Berlin, pp. 113-123.

Le Maitre, R.W., Bateman, P., Dudek, A., Keller, J., Lameyre, J., Le Bas, M.J., Sabine, P., Schmid, R., Sorensen, H., Streckeisen, A., Woolley, A.R., and Zanettin, B., (1989): *A Classification of Igneous Rocks and Glossary of Terms. Recommendations of the International Union of Geological Sciences Subcommision of the Systematics of Igneous Rocks*. Blackwell, London. 193pp.

Lottermoser, B.G., Ashley, P.M., (2000): Geochemistry, petrology and origin of Neoproterozoic ironstones in the eastern part of the Adelaide Geosyncline, South Australia. *Precambrian Research*, 101, pp. 49-67.

Martin, D., McB., Clendenin, C.W., Drapez, B., McNaughton, N.J., (1998): Tectonic and geochronological constraints on late Archaean and Palaeoproterozoic stratigraphic correlation within and between the Kaapvaal and Pilbara Cratons. *Journal of the Geological Society*, London 155, pp. 311-322.

McCrea, J.M., (1950): On the isotopic chemistry of carbonates and a paleotemperature scale. *Journal of Chemistry and Physics*, 18, pp. 849-857.

McLennan, S.M., (1989): Rare earth elements in sedimentary rocks: influence of provenance and sedimentary processes. In: Lipin, B.R and McKay, G.A., (Eds.), *Geochemistry and mineralogy of rare earth elements. Reviews in Mineralogy*, 21, pp. 169-200.

Moen, H.F.G., (1999): The Kheis Tectonic Subprovince, southern Africa: A lithostratigraphic perspective. *South African Journal of Geology*, 102, pp. 27-42

Netshiozwi, S.T., (2002): *Origin of high-grade hematite ores at Thabazimbi mine, Limpopo Province, South Africa*. Unpublished, M.Sc. thesis, Rand Afrikaans University, Johannesburg, 135 pp.

Pilipiuk, A.N., Ivanikov, V.V., Bulakh, A.G., (2001): Unusual rocks and mineralization in a new carbonatite complex at Kandaguba, Kola Peninsula, Russia. *Lithos*, 56, pp. 333-347.

Pineau, F., Javoy, M., and Allegre, C.J., (1973): Etude systematique des isotopes de l'oxygene, du carbone et du strontium dans les carbonatites. *Geochimica et Cosmochimica Acta*, 37, pp 2363-2377.

Reichardt, F.J., (1994): The Molopo Farms Complex, Botswana: history, stratigraphy, petrography, petrochemistry and Ni-Cu-PGE mineralization. *Exploration and Mining Geology* 3, pp. 264-284.

Reid, D.L., (1982): Age relationship within the Vioolsdrif batholith, lower Orange River region. II. Two stage emplacement history and the extent of Kibaran overprinting. *Trans. Geol. Soc. S. Afr.* 85. pp. 105-110.

Reid, D.L., (1997): Sm-Nd age and REE geochemistry of Proterozoic arc-related igneous rocks in the Richtersveld Subprovince, Namaqua Mobile belt, Southern Africa. *Journal of African Earth Science*. 24. pp. 621-633.

Rollinson, H. (1993): *Using geochemical data: evaluation, presentation, interpretation*. Longman Group Ltd., Prentice Hall, Essex.

S.A.C.S. (South African Committee for Stratigraphy), (1980): Stratigraphy of South Africa. Part I (comp. Kent, L.E.). Lithostratigraphy of Republic of South Africa, South West Africa/Namibia, Transkei and Venda: Handbook. Geological Survey of South Africa, 8, 690 pp.

Schürmann, L.W., Horstmann, U.E., and Cloete, H.C.C., (1997): Geochemical and stable isotope patterns in altered volcanoclastic and intrusive rocks of the Kruidfontein Carbonatite Complex, South Africa. *Journal of African Earth Sciences*, 25. pp. 77-101.

Sumner, D.Y., Bowring, S.A., (1996): U-Pb geochronologic constraints on deposition of the Campbellrand Subgroup, Transvaal Supergroup, South Africa. *Precambrian Research*, 79, pp. 25-35.

Taylor, N., (1979): The stratigraphy of the early-Proterozoic Buffalo Springs Group in the Thabazimbi area, west-central Transvaal. *Transaction Geological Society of South Africa* 82, pp. 215-226.

Taylor, H.P.J., Frechen, J., and Degens, E.T., (1967): Oxygen and carbon isotope studies of carbonatites from the Laacher See District, West Germany and the Alnö District, Sweden. *Geochimica et Cosmochimica Acta*, 31, pp. 407-430.

Taylor, S.R., McLennan, S.M., (1981): The composition and evolution of the continental crust: rare earth elements evidence from sedimentary rocks. *Phil. Trans. R. Soc.*, A301, pp. 381-399.

Taylor, S.R., McLennan, S.M., (1985): *The continental crust: its composition and evolution*. Blackwell, Oxford.

Taylor, D., Dalstra, H.J., Harding, A.E., Broadbent, G.C., and Barley, M.E., (2001): Genesis of high-grade hematite orebodies of the Hamersley Province, Western Australia, *Economic Geology*, 96. pp. 837-878.

Thomas, R.J., Cornell, D.H., Moore, J.M., and Jacobs, J., (1994): Crustal evolution of the Namaqua-Natal Metamorphic Province, southern Africa. *South African Journal of Geology*, 97(1), pp. 8-14.

Vajner, V., (1974): The Doringberg Fault and its relation to the post-Waterberg deformation. *Transvaal Geological Society of South Africa*. 77, pp 295-300.

Miyano, T., and Beukes, N.J., (1984): Phase relations of stilpnomelane, ferri-annite, and riebeckite in very low-grade metamorphosed iron-formations. *Transvaal Geological Society of South Africa*. 87, pp. 111-124.

Morris, R.C., and Horwitz, R.C., (1983): The origin of the Iron-Formation-rich Hamersley group of Western Australia – deposition on a platform. *Precambrian Research*, 21, pp. 273-297.

Van Deventer, J.L., Eriksson, P.G., and Snyman, C.P., (1986): The Thabazimbi iron ore deposits, north-western Transvaal. In: Mineral Deposits of Southern Africa. Vols. I & II. *Geol.Soc.S.Afr.*, Johannesburg.

Van Niekerk, H.S., (1998): *The Paleoproterozoic Olifantshoek-Kheis orogen and its mineral potential*. Unpublished. Progress Report, Johannesburg, pp. 1-26.

Van Niekerk, H.S., Gutzmer, J., Beukes, N.J., Smit, A., Armstrong, R., Yaspakurt, V., and Vasconcelos, P., (2003): *The Grenvillian assembly of Rodinia: Timing of accretion on the western margin of the Kalahari (KAAPVAAL) craton*. Unpublished report. Rand Afrikaans University, Johannesburg, pp. 22.

Van Wyk, J.P., (1980): *Die geologie van die gebied Rooinekke-Matsap-Wolhaarkop in Noord-Kaapland met spesiale verwysing na die Koegas-Subgroep, Transvaal-Supergroep*. Unpublished. M.Sc. thesis, Rand Afrikaans University, Johannesburg, 159pp.

Walter, A.V., Flicoteaux, R., Parron, C., Loubet, M., Nahon, D., (1995): Rare-earth elements and isotopes (Sr, Nd, O, C) in minerals from the Juquiá carbonatite (Brazil): tracers of a multistage evolution. *Chemical Geology*, 120, pp. 27-44.

Webb, A.D., Dickens, G.R., Oliver, N.H.S., (2003): From banded iron-formation to iron ore: geochemical and mineralogical constraints from across the Hamersley Province, Western Australia. *Chemical Geology, in press*, 14123, pp. 1-37.

Wilson, M.G.C., Anhaeusser, C.R., (Editors) (1998): *The mineral resources of South Africa*. Council of Geoscience, 6th Edition.

Winterbach, J.C., (1979): *Nauga Hematite Ore Deposit*. Unpublished Rand Mines, Limited report. 3pp.

Woolley, A.R., and Kempe, D.R.C., (1989): Carbonatites: nomenclature, average chemical composition, and element distribution. In: *Carbonatites: Genesis and Evolution* (eds) Bell, K., pp. 1-14. Unwin Hyman, London.

