

LISTS OF FIGURES

Pages

- Fig 1-1:** Simplified geological map of the Precambrian of Madagascar adapted from Tucker et al. (1998), showing the area of study within the Precambrian of Madagascar (Le Grange, 2001). 2
- Fig 2-1:** 1/200,000,000 Metallogenic map of Madagascar modified after Razafinimparany (1978), showing gold occurrences throughout the Malagasy Precambrian. 5
- Fig 2-1:** Geological map of the study area, with sample localities. Modified from the published 1/100,000 geological maps of Alsac (1963) and Moine (1963). 9
- Fig 2-2:** Photograph of a strongly foliated amphibolite outcrop forming a riverbed. The foliation strikes NW-SE and dips 20° SW. Locality 162, GPS: 19° 35' 57.2" S, 45° 33' 19.7" E. 10
- Fig 2-3:** Photograph showing mm-thick felsic veins oriented parallel to the fabric of the amphibolite. The amphibolite has a speckled appearance due to 1-2 mm amphibole grains in rosette shape, which is shown by the small white dots on the photograph. Locality 191, GPS: 19° 39' 01.0" S, 45° 44' 33.9" E. 10
- Fig 2-4:** Photograph showing the sequence of quartzofeldspathic and biotite-rich layers of the quartzofeldspathic gneisses, with predominance of the felsic layers. Locality 91, GPS: 19° 37' 17" S, 45° 38' 28" E. 11
- Fig 2-5:** a) Photograph showing pegmatite boudins within the felsic layer of the quartzofeldspathic gneiss. The boudins are oriented parallel to the fabric of the rock. Locality 163, GPS: 19° 37' 05.5" S, 45° 38' 25.0" E. b) Photograph showing recumbent folding of the quartzofeldspathic layer. The fold plane strikes 145° and dips 6° W, and the fold axis is almost horizontal, trending 175°. Locality 69, GPS: 19° 37' 31" S, 45° 41' 02" 12
- Fig 2-6:** Photograph showing recumbent folding of the quartzofeldspathic layer. The fold plane strikes 145° and dips 6° W, and the fold axis is almost horizontal, trending 175°. Locality 69, GPS: 19° 37' 31" S, 45° 41' 02" E. 12
- Fig 2-7:** Photograph showing an outcrop of marble and calc-silicate layer (1-2 m) within the amphibolite. The exposed surface of the carbonate unit is parallel to the foliation. Locality 162, GPS: 19° 35' 57.2" S, 45° 33' 19.7" E. 13
- Fig 2-8:** Photograph of a 3-4 m wide mafic body within the carbonate unit. Locality 126, GPS: 19° 33' 28.6" S, 45° 38' 04.7" E. 13
- Fig 2-9:** Photograph of a foliated granodiorite from the Dabolava pluton, with coarse plagioclase up to 1 mm across, forming augen texture. Locality 51, GPS: 19° 38' 12" S, 45° 44' 00" E. 15
- Fig 2-10:** Photograph showing a foliated granodiorite rock from the Ambatomiefy 15

pluton, with a 50 cm thick fine-grained mafic enclave, which is oriented parallel to the fabric of the host rock. Granodiorite xenoliths are observed within the mafic body. Locality 85A, GPS: 19° 35' 03" S, 45° 39' 02" E.

Fig 2-11: Photograph showing a 30 cm thick granodiorite dyke within the amphibolite. Visible within the dyke are xenoliths of the amphibolite. Locality 136, GPS: 19° 33' 01.0" S, 45° 37' 31.4" E. 16

Fig 2-12: Photograph of an outcrop of amphibolite crosscut by a weathered granodioritic dyke, 60cm thick. The amphibolite contains numerous randomly oriented mm-thick felsic veins. Locality 124, GPS: 19° 33' 24.8" S, 45° 37' 44.7" E. 17

Fig 2-13: Photograph showing an outcrop of a foliated granodiorite rock from Andasibe pluton, with mafic slivers, crosscut by a 50 cm wide amphibolite dyke, trending 325°. The amphibolite presents the same characteristic observed elsewhere, with numerous mm-thick felsic veins. Locality 136, GPS: 19° 33' 01.0" S, 45° 37' 31.4" E. 17

Fig 2-14: Photograph showing an outcrop of deformed granodiorite rocks from the Ambatomiefy pluton. The deformation is defined by a folded pegmatite. A later tourmaline-bearing pegmatite dyke crosscuts the fabric of the host rock. Locality 122, GPS: 19° 34' 29."9 S, 45° 37' 23.5" E. 18

Fig 2-15: Photograph showing granodiorite dykes crosscutting the gabbro. The dykes present joints that are parallel to the foliation of the gabbro. Locality 77, GPS: 19° 38' 14" S, 45° 40' 11" E. 19

Fig 2-16: Photograph showing gabbro xenoliths within the granodiorite. Locality 77, GPS: 19° 38' 14" S, 45° 40' 11" E. 19

Fig 2-17: Photograph showing an outcrop of weathered gabbro, with composition layering. Locality 08, GPS: 19° 38' 28" S, 45° 47' 59" E. 20

Fig 2-18: Geological map of the area of study showing trend of the gold-bearing quartz veins and the orientations of the foliation of the host granodiorite plutons and the surrounding country rocks, with the trend of the fabrics of the rocks according to aerial photographs interpretations, and field measurements. Modified from Alsac (1963) and Moine (1963). 22

Fig 2-19: Photograph of an outcrop of folded amphibolite, showing a minor anticline. Locality 29, GPS: 19° 38' 56" S, 45° 45' 04" E. 23

Fig 2-20: Photograph of a complex structure found within the quartzofeldspathic gneisses, with minor amphibolitic layers. A resistant pegmatite layer, consisting mainly of quartz and albite, is deformed with the rock package and forms boudins. 24

Locality 88, GPS: 19° 37' 09" S, 45° 38' 58" E.

- Fig 2-21:** Photograph of a 3-4m wide E-W trending shear zone within the amphibolites that contain 10-30cm thick tonalitic layers. The deformation is crosscut by a later N-S trending foliated amphibolite dyke. 24
- Fig 2-22:** U-Pb isochron diagram for granodiorite sample from the Ambatomiefy pluton (RRH 98-131F), showing an upper intercept age of 1002 ± 2 Ma. 26
- Fig 2-23:** U-Pb isochron diagram for a gabbro sample from the area of study, with an upper intercept age of 982 ± 2 Ma. 27
- Fig 3-1:** Photomicrograph of pyroxene-bearing amphibolite (sample RRH 28), with coarse orthopyroxene grains and finer grained amphiboles, in a groundmass consisting of plagioclase with minor quartz. Taken in plane polarized light. 29
- Fig 3-2:** Photomicrograph of amphibolite sample (RRH 45) taken in plane polarized light and showing hornblende rimming clinopyroxene. 29
- Fig 3-3:** Photomicrograph of amphibolite sample RRH 72, taken in crossed polarizers, showing epidote grains in a groundmass of microcline, quartz and plagioclase. 30
- Fig 3-4:** Modal Plag-Hb-Opx+Cpx ternary diagram of the amphibolites showing the variation in their mineralogy. 31
- Fig 3-5:** Photomicrograph of the quartzofeldspathic layer (sample RRH 27), taken in crossed polarizers, showing subhedral coarse muscovite grains defining the foliation of the rock with brown biotites. 32
- Fig 3-6:** Photomicrograph of a biotite-rich layer of the quartzofeldspathic gneisses, with biotite defining the foliation of the rock (sample RRH 174), and showing equant garnet. 33
- Fig 3-7:** Modal QKP diagram of the quartzofeldspathic gneisses, the granodiorite plutons and the granitic intrusions. 34
- Fig 3-8:** Photomicrograph of sample RRH 181A, a sillimanite-bearing feldspathic quartzite, showing elongated sillimanite defining the foliation, within a groundmass consisting of plagioclase and quartz. Taken in crossed polarizers. 35
- Fig 3-9:** Photomicrograph of a rock that forms the impure layer within the marbles (sample RRH 62). It shows epidote and hornblende grains in a groundmass of calcite with minor quartz and plagioclase. Taken in plane polarized light. 36
- Fig 3-10:** Photomicrograph of a granodiorite sample from the Ambatomiefy pluton (RRH 136), with biotite defining the foliation of the rock. Taken in crossed polarizers. 37

- Fig 3-11:** Photomicrograph of a gabbro sample (RRH 104) taken in plane polarized light showing hornblende mantling clinopyroxene within a plagioclase and quartz groundmass. 38
- Fig 3-12:** Photomicrograph of a foliated granite with microcline phenocrysts in a fine grained felsic ground mass. Biotites define the foliation of the rock (sample RRH 117). Taken under crossed nicols. 39
- Fig 4-1:** Jensen diagram classifying the amphibolites according to their cation percentages of Al, (Fe_(total) + Ti), and Mg. According to the diagram, the composition of the amphibolites consists mainly of tholeiite basalts with minor andesite and basalt. The boundary line between the tholeiitic and calc-alkaline fields is defined by Rickwood (1989) (Rollinson, 1993). 41
- Fig 4-2:** AFM diagram for the amphibolites showing tholeiitic affinities. The line separating the fields of the tholeiitic and calc-alkaline series is from Irvine and Baragar (1971). 42
- Fig 4-3:** a) P₂O₅-Zr diagram for the amphibolites to determine the affinities of their magma origin. b) TiO₂-Zr/P₂O₅ diagram, same purpose as the former diagram (Floyd and Winchester, 1976; Rollinson, 1993). 43
- Fig 4-4:** Zr/Y-Ti/Y diagram discriminating the within-plate basalt from the other basalt types. The boundary line between the two fields is from Pearce and Gale, 1977. Data points show compositions of amphibolites from the Dabolava area. 44
- Fig 4-5:** Ti-Zr-Y discrimination diagram for basalts containing four fields which consist of field A for island-arc tholeiites, field C for calc-alkali basalts, field D for within-plate basalts and field B for MORB, island-arc tholeiites and calc-alkali basalts. (Pearce and Cann, 1973). Data points show compositions of amphibolites from the Dabolava area. 45
- Fig 4-6:** Diagram that subdivides continental and oceanic arcs; the boundary line is based on a Zr/Y value of 3 (Rollinson, 1993). 45
- Fig 4-7:** DF values vs. SiO₂ concentration for the quartzofeldspathic gneisses, showing a dominant igneous parentage for the rocks. 47
- Fig 4-8:** Shand's index for the quartzofeldspathic gneisses. A=Al₂O₃, C=CaO, N=Na₂O, K=K₂O 47
- Fig 4-9:** Total alkalis-silica diagram (TAS) for the quartzofeldspathic gneisses. The analyses were normalized 100wt% anhydrous. Q = normative quartz, Ol = normative olivine. Rollinson, 1993. 48
- Fig 4-10:** Harker diagrams for quartzofeldspathic gneisses. 49

Fig 4-11: Trace elements concentration (ppm) vs. SiO ₂ (wt%) for the quartzofeldspathic gneisses.	51
Fig 4-12: Normative QKP diagram to classify the plutons.	52
Fig 4-13: Shand's index for the granodiorite plutons. A=Al ₂ O ₃ , C=CaO, N=Na ₂ O, K=K ₂ O.	52
Fig 4-14: SiO ₂ vs A/CNK diagram showing the subdivision according to Shand's index and classification of the granitoids according to their presumed origin (I-, S-type) (Chappell and White, 1974; Clarke 1992).	53
Fig 4-15: Harker variation diagrams for the granodiorite plutons, plotting oxides (Al ₂ O ₃ , Fe ₂ O ₃ , MgO, TiO ₂ , P ₂ O ₅ , CaO, MnO, Na ₂ O, K ₂ O) against SiO ₂ .	54
Fig 4-16: Trace elements concentration (ppm) vs. SiO ₂ (wt%) for the granodiorite plutons.	56
Fig 5-1: Photograph of an excavation by local artisans, within the Dabolava pluton, showing the gold-bearing quartz vein hosted by granodiorite rocks.	59
Fig 5-2: Close-up of the gold-bearing quartz veins, which form multiple layers with thickness varying from 2-5cm.	60
Fig 5-3: a) Photomicrograph of the gold-bearing quartz vein from the Ambatomiefy pluton, showing gold and pyrrhotite associated with the hydrothermal biotite. b) Photomicrograph showing pyrrhotite overgrowing gold and both minerals are associated with biotite. The width of the view is approximately 2 mm across, and the photomicrographs are taken in reflected plane polarized light.	62
Fig 5-4: Ternary diagram showing compositions of the gold grains, which mainly consists of 94-97% gold.	63
Fig 5-5: Geological map of the area of study showing the trend of the gold-bearing quartz veins and the orientations of the foliation of the host granodiorite plutons and the surrounding country rocks, with the trend of the fabrics of the rocks according to aerial photographs interpretations. Modified from Alsac (1963) and Moine (1963).	65
Fig 5-6: Photomicrograph of fluid inclusions in quartz grain from the sample RRH 196D. Type 1-b and type 2 inclusions form scattered groups. The width of the view is 0.5 mm.	68
Fig 5-3: a) Histograms representing the melting temperature of the carbonic phase (T _m CO ₂) for type 2 (in red) and type 1-b (in blue) fluid inclusions. b) Temperature of melting of clathrate (T _m clath) for type 2 fluid inclusions. c) Histograms of the homogenization temperature of carbon dioxide, into the liquid or vapor phase,	66

$T_h\text{CO}_2$ (L) and $T_h\text{CO}_2$ (V), respectively, for type 2 and type 1-b inclusions. d) Homogenization temperatures and decrepitation temperatures (T_d) for the two types of fluids.

Fig 5-8: a) Histograms representing the melting temperature of the carbonic phase ($T_m\text{CO}_2$) for type 2 (in red) and type 1-b (in blue) fluid inclusions. b) Temperature of melting of clathrate ($T_{m\text{clath}}$) for type 2 fluid inclusions. c) Histograms of the homogenization temperature of carbon dioxide, into the liquid or vapor phase, $T_h\text{CO}_2$ (L) and $T_h\text{CO}_2$ (V), respectively, for type 2 and type 1-b inclusions. d) Homogenization temperatures and decrepitation temperatures (T_d) for the two types of fluids. 70

Fig 5-9: Raman spectrum of the CO_2 -rich- H_2O - NaCl inclusion (type 2) showing the strongly pronounced CO_2 peaks (b), the N_2 and CH_4 peaks, which are at 2331 Rcm^{-1} and 2917 Rcm^{-1} respectively are absent on the spectrum (c, d). 71

Fig 5-10: Raman spectrum of the two-phase CO_2 -rich inclusion (type 1-b) with the two CO_2 peaks at 1285 and 1388 Rcm^{-1} . The N_2 and CH_4 peaks are absent (Fig 5-5c, d). 72

LIST OF TABLES:

	Pages
Table 2-1: Isotope dilution analyses of zircons from the granodiorite sample RRH98 131F, from the Ambatomiefy pluton.	26
Table 4-2: DF values for the quartzofeldspathic gneisses.	46
Table 5-1: Description of the characteristics of the gold-bearing quartz veins encountered in the different gold prospects found within the Dabolava and the Ambatomiefy plutons, with their structural orientations.	60
Table 5-2: Microprobe analyses of gold grains showing the concentrations of the different elements within the gold. Gold grains are from gold-bearing quartz veins hosted by the Dabolava granodiorite pluton.	64
Table 3-1: Modal proportions of rock-forming minerals within the different lithologies studied.	Appendix
Table 4-1: Major and trace element analyses of the amphibolites, quartzofeldspathic gneisses, granodiorite plutons and granitoid bodies, from the area of study, with their CIPW Norm calculations.	Appendix
Table 5-1: Microthermometry results of fluid inclusion studies of a sample of gold-bearing quartz vein (RRH 196) from the Ambatomiefy pluton.	Appendix