

Low Intensity Laser Irradiation at 636 nm Induces Increased Viability and Proliferation in Isolated Lung Cancer Stem Cells

Anine Crous, MTech, and Heidi Abrahamse, PhD

Abstract

Objective: The purpose of this *in vitro* study was to evaluate the effects of low-intensity laser irradiation (LILI) on isolated lung cancer stem cells (CSCs) after several time intervals, using a wavelength of 636 nm and fluences between 5 and 20 J/cm². **Background data:** LILI has been proven to have a biomodulatory effect on various diseased conditions. A number of studies have been conducted on CSCs. **Methods:** Lung CSCs were isolated from lung cancer cells (A549), using cell surface marker CD 133. Isolated lung CSCs were divided into four groups: group 1 consisted of control cells receiving no irradiation; groups 2, 3, and 4 were exposed to laser irradiation at fluences of 5, 10, and 20 J/cm², respectively. LILI was performed using a 636 nm diode laser with a power output of ±85 mW. Cellular responses were evaluated after 24, 48, or 72 h, and included cell morphology, viability, and proliferation. **Results:** Cellular morphology indicated an increase in cell density caused by cell proliferation over time. Biostimulatory effects were achieved in lung CSCs when examining viability and proliferation. **Conclusions:** It should, therefore, be noted that a low wavelength of 636 nm at various fluences induces biostimulation, which may have detrimental effects when using LILI as a form of regeneration.

Introduction

LUNG CANCER ACCOUNTS FOR MORE cancer-related mortalities worldwide than any other cancer in both men and women¹ Treatments for lung cancer include chemotherapy, radiation, and surgery. Despite these therapeutic advances, lung cancer still has a poor prognostic outcome, which can be attributed to its metastatic potential.² It has been suggested that lung cancer can be driven by a small subpopulation of self-renewing cells that can sustain malignant growth, called “cancer stem cells” (CSCs) or “tumor-initiating cells.” These cells share various characteristics with normal stem cells (SC) such as self-renewal and multipotent differentiation.³ An important difference between normal SCs and CSCs, which researchers are proposing, is their dependability on their stem cell niche. A stem cell niche is a specialized microenvironment with the following features and functions: it comprises of a group of cells situated in a special location that maintains the SCs, it functions as a physical anchoring site between the SCs and their environment, it generates extrinsic factors that control SC proliferation and lineage fate, and it also controls the asymmetric division of these cells.⁴ CSCs evade this advanced niche, signalling control quiescence and proliferation as a result of its genetic mutation, making it

self-sufficient. It has been reported that deregulation of the SC niche leading to uncontrolled proliferation of SCs can lead to tumorigenesis.⁵ CSCs have also been found capable of metastasizing, being resistant to current treatment methodologies such as chemo- and radiation therapy, and being resistant to cell death.⁶ An explanation for why chemotherapy fails to eradicate CSCs is its capability to only remove rapidly proliferating cells, and CSCs are usually slow in multiplying and, therefore, unresponsive to this treatment. As a result of current cancer treatments failing, new treatment methods that will have a good prognosis, and fewer side effects are being investigated. One of the current therapies under investigation is photodynamic therapy (PDT), in which low-level light-emitting lasers are used to activate a photochemotherapeutic compound called a “photosensitizer.”

Low-intensity laser irradiation (LILI) is a photobiomodulative therapy that makes use of low intensities of light that is emitted coherently at a specific wavelength in the visible spectrum, which consists of red and near-infrared (IR) light between 600 and 1070 nm, also known as the optical window where effective tissue penetration is maximal. This therapeutic model exposes cells or tissues to visible red light and near-IR light, having either a biostimulatory or a bioinhibitory effect.⁷ The mechanism by

which LILI induces a photobiological effect is chromophore stimulation at the mitochondrial level, causing various metabolic effects, depending upon the wavelength and fluence used.⁸

Previous studies conducted using LILI with wavelengths between 625 and 675 nm and fluences ranging from 1 to 15 J/cm² all indicated a biostimulatory effect showing an increase in proliferation and viability on various cells;^{9–13} however, studies indicated biochemical inhibition when using higher wavelengths of 800–830 nm and fluences ≥ 10 J/cm².^{2,10,14,15}

Photobiomodulation relies on specific parameters such as wavelength, fluence, power density, pulse structure, and treatment time when applied to biological tissue. This allows for targeting of specific light-absorbing molecules in specific tissues, operating on the principle of photochemistry, as opposed to photothermogenesis. The light energy absorbed causes singlet state excitation of oxygen molecules, leading to triplet state excitation causing an energy transfer to ground state molecular oxygen (a triplet) to form the reactive species, singlet oxygen. Alternatively superoxide may be formed as a result of electron reduction. LILI operates at an exact wavelength of light, which influences the depth of tissue penetration. Similar to normal cells, cancer cells also contain with intracellular chromophores. Different cellular chromophores are stimulated at different wavelengths.^{16,17} Therefore, the prediction can be made that in targeting cancerous cells, the outcome expected can be controlled by the wavelength as well as by the energy output that will lead to either stimulation or inhibition. The exact mechanism behind the stimulation of the light-absorbing molecules producing these two different effects is still being investigated.

CSCs can be identified and isolated using their antigenic markers.¹⁸ Prominin 1 (CD 133) is a pentaspan transmembrane glycoprotein usually found on cell surfaces.¹⁹ It has been used to identify and isolate different SCs and CSCs.²⁰ Lung CSCs have previously been isolated using the surface marker CD133,²¹ and the same antigenic marker was used in this study. Control cell lines used were the CaCo-2 (Caco-2, HTB-37™) cell line, which is a continuous cell of heterogeneous human epithelial colorectal adenocarcinoma cells, which have been found to express CD 133.²⁰ SKUT-1 (SKUT-1, HTB114™) is a cell line derived from human uterus leiomyosarcoma cells.²² These cells do not express the surface marker CD 133, as they were used as a somatic cell line and contributed as the negative control cell line. In this study, we aimed to evaluate the biomodulative effects induced by LILI on isolated lung CSCs. It resulted that there were possible detrimental effects associated with LILI when it was used in tissue with underlying malignancy, as LILI provides effective tissue penetration and can produce similar results to the cell layer, as was found in this study.

Materials and Methods

Cell culture

This study used commercially obtained lung cancer cells (A549, ATCC[®] CCL-185). Rosewell Park Memorial Institute 1640 medium (RPMI) (Sigma, R8758), supplemented with 10% fetal bovine serum (FBS) (Biobrom, S0615), 0.5% penicillin/streptomycin (Sigma, P4333), and 0.5%

amphotericin B (Sigma, A2942) was used to culture these cells in. The cells were incubated at 37°C in 5% CO₂ and 85% humidity. Cells were grown to confluence before isolation of the CSCs. Upon lung cancer cells reaching confluence, the cells were washed twice with Hank's balanced salt solution (HBSS) (Sigma, H9269) and detached from the culture dish using 1 mL/25 cm² of TrypLE Express (Gibco, 12604).²³ For experimental purposes, isolated lung CSCs were cultured in complete medium using only 5% FBS. Cells were seeded at a final concentration of 2×10^4 cells in 3 mL complete medium, and cultured in a 3.4 cm² diameter culture dish followed by incubation for 24 h to allow cells to attach.

Control cell lines were cultured in Dulbecco's Modified Eagle's Media (DMEM) (Sigma, D5796) supplemented with 10% FBS, 2 mM L-Glutamine (Sigma, G7513), 1% penicillin/streptomycin (Sigma, P4333), and 1% amphotericin B (Sigma, A2942) for CaCo2 (ATCC[®], HTB-37™) which was used as the positive control,² and DMEM (Sigma, D5796) supplemented with 10% FBS (Biobrom, S0615), 1% penicillin/streptomycin (Sigma, P4333), and 1% amphotericin B (Sigma, A2942) for SKUT-1 (ATCC[®], HTB114™) as the negative control.²⁴

Isolation of lung CSCs

Lung CSCs were isolated using the magnetic bead isolation kit (Miltenyi Biotec, QuadroMACS™ separation unit 130-091-051), where they were magnetically labelled with microbead conjugated antibodies directed at the antigenic surface marker of interest. Lung CSCs were enriched using the CD133 MicroBead Kit (Miltenyi Biotec, CD133 MicroBead Kit, human 130-050-801) designed for the positive selection of cells expressing human CD133 antigen.

A single cell suspension was prepared, where the cell number was determined. The cells were spun down and resuspended in 80 μ L buffer per 10^7 total cells, and 20 μ L MicroBeads per 10^7 total cells were added and incubated for 15 min at 2°–8°C. Cells were washed, centrifuged, and resuspended in 500 μ L buffer per 10^8 cells. The separation column was prepared by running buffer through and discarding the effluent. The cell suspension was applied to the prepared column, where the unlabelled cells were collected and discarded and the positively selected cells in the column were flushed out into a suitable collection tube using buffer and a plunger.²⁵

Immunofluorescence

Fluorescent labelling of surface antigens with CD 133 (Prominin-1), which is a monoclonal antibody hosted in mouse and reacting in humans (USBio, 030034) on isolated cells, was used to confirm whether the lung CSCs that were isolated were of CSC origin.²⁶ Isolated cells along with control cell lines CaCo2 and SKUT-1 were cultured on heat-sterilized cover slips placed in Petri dishes (3.5 cm diameter), at a concentration of 2×10^5 cells in complete media. Cells were incubated overnight to attach to the cover slip. Cultures were removed from the incubator and washed twice with ice-cold phosphate-buffered saline (PBS)/bovine serum albumin (BSA)/azide buffer (PBS, Sigma, P4417), 0.1% w/v BSA (Sigma, A2153), and 0.01% w/v azide (Sigma, S8032) on ice, and then incubated with 10% serum

(Normal Goat Serum, abcam, ab7481) for 30 min on ice as a blocking step, and then washed twice again. Cells were then incubated with 100 μ L primary antibody rabbit anti-human CD 133⁺ (Abnova, PAB12663) in working buffer (2% serum in PBS BSA/azide buffer) for 30 min on ice. Cells were then rinsed three times with PBS BSA/azide buffer and incubated with 100 μ L of the secondary fluorescent antibody (abcam, ab6717) in working buffer for 30 min on ice, protected from light. Cells were rinsed three times as before and fixed in 4% paraformaldehyde for 10 min. After fixation, cells were rinsed once briefly with PBS, and then once with tap water before being stained with 4'-6-diamidino-2-phenylindole (DAPI) (Sigma, D9564) and mounted on glass slides using 0.1 M propylgallate (Sigma, 02370). Slides were viewed using a fluorescent microscope (Carl Zeiss, Axio Observer Z1).¹⁴

Laser irradiation

After 24 h of incubation, the culture medium was removed and the cells were washed with HBSS. Fresh medium was added, and cells were irradiated using a 636 nm diode laser (LTIA000-PLT20-636nm) provided by the National Laser Center (NLC) of South Africa. The power output was measured using the FieldMate Laser Power Meter (Detector S/N: 125J07R) and the value was used to calculate the exposure time. Cells were irradiated with energy densities of 5, 10, and 20 J/cm². Cells were irradiated from above, at room temperature with the culture dish lid off. Irradiation was performed in the dark, omitting nuisance variables that would interfere with the laser effect, such as polychromatic light. Laser parameters are shown in Table 1. Cell cultures were divided into four study groups. Group 1 was an unirradiated control, group 2 was irradiated at 5 J/cm², group 3 was irradiated at 10 J/cm², and group 4 was irradiated at 20 J/cm². Post-irradiation, cells were incubated for 24, 48, or 72 h.

Cell morphology

Morphological changes in the four different study groups of isolated lung CSCs were observed and studied using an inverted light microscope (Wirsam, Olympus CKX41) 24, 48, or 72 h post-irradiation. Morphological pictures were taken with the SC30 Olympus camera.

Adenosine triphosphate (ATP) cell viability assay

The number of metabolically active cells was determined using the CellTiter-Glo[®] luminescent cell viability assay

TABLE 1. LASER PARAMETERS USING THE 636 NM DIODE LASER

Parameters	Value
Laser type	Semiconductor (diode)
Wavelength (nm)	636
Wave emission	Continuous wave
Power output (mW)	85
Power density (mW/cm ²)	9.36
Spot size (cm ²)	9.1
Fluence (J/cm ²)	5, 10, and 20
Duration of irradiation \pm min, sec	8 min 54 sec, 17 min 48 sec, 35 min 36 sec

(Whitehead Scientific, Promega, G7573). This homogeneous method determines the number of viable cells in culture based on ATP quantitation. The assay utilizes the properties of a proprietary thermostable luciferase, which generates a luminescent signal proportional to the amount of ATP present released upon cell lysis. According to the manufacturer's protocol, 50 μ L of reconstituted reagent was added to an equal volume of cell suspension. The contents were mixed in a shaker for 2 min to induce cell lysis. The contents were then incubated at room temperature for 10 min to stabilize the luminescent signal.²⁷ The amount of ATP was quantified, and luminescence was recorded using the Perkin Elmer, VICTOR3[™] Multilabel Counter (model 1420) in relative light units (RLUs).

Trypan blue exclusion assay

Trypan blue staining method was used to quantify percentage viable cells. The assay is employed to identify dead cells, as cells that are viable have intact membranes and can effectively exclude the dye, whereas dead cells with compromised membranes become stained. Equal volumes of 0.4% Trypan blue (Invitrogen[™], Trypan Blue Stain [0.4%] T10282) and cell suspension were mixed and loaded into a counting chamber, where the number of cells that were viable and dead was counted using the Countess[®] Automated Cell Counter. Percentage viability was determined by calculating the number of viable cells from the total number of cells.

Alamar blue cell proliferation assay

Alamar blue is a quantitative assay that uses a fluorometric growth indicator based on detection of metabolic activity. It uses a redox indicator, where metabolically active cells will reduce the dye Resazurin, which is blue, to its reduced form Resorufin, which is a fluorescent red. Then, 100 μ L of cell suspension was added to a 96 well microtiter plate, and 10 mL AlamarBlue[®] reagent (Invitrogen[™], AlamarBlue[®] DAL1025) was added. The plate was incubated for 3 h (37°C in 5% CO₂). Fluorescence was then measured using the Victor-3 (Perkin-Elmer, Separation Scientific) at Ex/Em 560/590.

Statistical analysis

The isolated lung CSC cell lines were used between passage numbers one and four. Experiments were repeated four times ($n=4$). All assays were performed in duplicate, and the average was used. Results were statistically analysed using Sigma Plot Version 8.0, and the mean, standard deviation, and standard error were obtained. A student *t* test and one way ANOVA were performed to detect differences between the control and experiments and between experimental groups, as well as differences between controls and treated samples from 24 to 72 h. Statistical significances are indicated in the figures as $p < 0.05$ (*), $p < 0.01$ (**), and $p < 0.001$ (***)

Results

Immunofluorescence

Expression of the CD 133 stem cell marker after CSC isolation was determined by indirect immunofluorescence

TABLE 2. PERCENTAGE VIABILITY OF IRRADIATED AND CONTROL CELLS

Time	Fluence					
	5 J/cm ²		10 J/cm ²		20 J/cm ²	
24 h	64.5	±4.9075	71.25	±6.9537	70.625	±2.7186
48 h	88.325	±3.0682	69	±11.158	69.875	±6.9263
72 h	89.75	±4.2647	87	±2.7003	90.625	±2.3038
	92.625	±1.8861	91.75	±2.1747	84.875	±2.5362
	91.875	±1.519	92.375	±1.1434	95.375	±0.826
			95.625	±0.4732	93.375	±1.2479

Control ± SE.

Trypan blue exclusion (percentage) was performed to assess cellular viability of different irradiation fluences and measured after several time intervals. Differences were noted when comparing viability from 24 to 72 h post-irradiation, showing an increase in viability over time from all different fluences used.

cancer treatments, but also about LILI when used as a therapeutic model. Cancer treatments such as surgery, radiotherapy, and chemotherapy still offer restricted prognostic outcome when it comes to competence, and produce undesirable side effects. Therefore, alternate therapeutic options are essential, and need to be investigated.²³ The focus needs to be turned to CSC-targeted therapy. LILI on the other hand which has been used for its desired effects of increased cell proliferation and viability,⁹ can be harmful, bearing in mind CSC characteristics.

Previous studies conducted on lung cancer indicated that CD 133 positive cells displayed a greater ability for self-renewal, tumor initiation, and drug resistance.²⁹ In previous studies conducted using neuroblastoma (NB) tumor samples expressing CD133, the marker was shown to repress cell differentiation and accelerated cell proliferation, anchorage-independent colony formation, and *in vivo* tumor formation,³⁰ all indicative of CSC characteristics.

In this study, we explored the possibility of outcomes using LILI on lung CSCs. Lung CSCs were successfully isolated using the CD 133 cell surface marker. Control cells receiving no irradiation maintained their cell morphology, whereas there was an increase seen in viability and proliferation as a result of normal cell cycle maintenance. These results concur with those of Mvula et al.¹³ In this study, cells exposed to laser irradiation showed a similar trend compared with control cells, although it should be noted that when the test cells were compared with their unirradiated controls over a time period of 24–72 h, statistical differences were obtained, indicating a remarkable increase in viability and proliferation at the different fluences used, showing a significant stimulation in the cell cycle leading to an exponential rate in mitosis. There were changes in cell morphology, increased cell viability, and increased cell proliferation. These results can be explained by the following: previous studies suggest that LILI can stimulate




FIG. 4. The Alamar blue assay was used to measure cell proliferation. Increase in proliferation is seen in all the samples as time elapses from 24 to 72 h. Unirradiated cells' proliferation time was compared with the treated cells' proliferation time. Statistical significant differences were expressed as $p < 0.001$ (***).

cells with a wavelength of 636 nm and a low fluence of 5 J/cm².¹³ One SC characteristic is that they stay quiescent until regeneration of cells stimulates the proliferation via a specific signal, whereas CSCs proliferate uncontrollably regardless of any cell signalling pathway,³¹ allowing the unirradiated control CSCs and irradiated CSCs to proliferate at a similar rate. CSCs are capable of self-renewal opposing cell death,⁶ even when exposed to higher fluences of 20 J/cm², suggesting that irradiation at 5, 10, and 20 J/cm² with a wavelength of 636 nm has no damaging effect, but rather a proliferatory effect on lung CSCs. This is then in agreement with previous studies also indicating that parameters such as wavelength, fluence, and intensity play an important role in photobiomodulation of cellular metabolism,^{10,12-14} which shows that wavelengths between 600 and 800 nm with fluences of 5–20 J/cm² can induce biostimulation, whereas wavelengths of 800–1070 nm with fluences of 20–40 J/cm² may be able to induce bioinhibition.

Conclusions

This study indicates the possible detrimental effect that LILI may have when used as a biostimulatory therapy on the underlying tissue CSCs when considering the proliferation and viability induced using visible wavelengths. Therefore, LILI should be used with caution clinically, bearing in mind the risk of stimulating potential underlying CSCs, as LILI provides optimum penetration depth in tissue, reaching underlining cell layers, which may react similarly to in this study. When considering the use of LILI as a bioinhibitory treatment of human lung cancer, one needs to explore these effects on lung CSCs, using higher wavelengths such as IR, because visible red light initiates CSC proliferation and viability. This study focused on lung CSCs at 636 nm. Other CSCs may respond differently than lung CSCs as it has been indicated in several studies that cancer cell lines differ with respect to their response to photobiomodulation. We propose that additional research should be conducted to establish the effects of LILI on lung as well as additional CSCs, determining the effect of wavelength and fluence.

Acknowledgments

This work was conducted at the Laser Research Centre at the University of Johannesburg and was supported by the National Laser Centre of South Africa.

Author Disclosure Statement

No competing financial interests exist.

References

- American Cancer Society. Selected Cancers. Cancer Facts & Figures, 2012 Available at: <http://www.cancer.org/research/cancerfactsfigures/cancerfactsfigures/cancer-facts-figures-2012> (Last accessed June 12, 2015).
- Manoto SL, Sekhejane PR, Houreld NN, Abrahamse H. Localization and phototoxic effect of zinc sulphothalocyanine photosensitizer in human colon (DLD-1) and lung (A549) carcinoma cells (in vitro). *Photodiagnosis Photodyn Ther* 2012;9:52–59.
- Rivera C, Rivera S, Loriot Y, Vozenin MC, Deutsch E. Lung cancer stem cell: new insights on experimental models and preclinical data. *J Oncol* 2011;2011:549181.
- Li L, Neaves WB. Normal stem cells and cancer stem cells: the niche matters. *Cancer Res*. 2006;66:4553–4557.
- Chepko G, Slack R, Carbott D, Khan S, Steadman L, Dickson RB. Differential alteration of stem and other cell populations in ducts and lobules of TGF alpha and C-Myc transgenic mouse mammary epithelium. *Tissue Cell* 2005;37:393–412.
- Crocker AK, Allan AL. Cancer stem cells: implications for the progression and treatment of metastatic disease. *J Cell Mol Med* 2008;12:374–390.
- Huang YY, Chen A, Hamblin M. Low-level laser therapy: an emerging clinical paradigm. *SPIE Newsroom*. 2009 Available at: <http://spie.org/x35504.xml> (Last accessed March 12, 2015).
- Karu T. Photobiology of low-power laser effects. *Health Phys* 1989;56:691–704.
- de Villiers JA, Houreld NN, Abrahamse H. Influence of low intensity laser irradiation on isolated human adipose derived stem cells over 72 hours and their differentiation potential into smooth muscle cells using retinoic acid. *Stem Cell Rev Nov* 2011;7:869–882.
- Moore P, Ridgway TD, Higbee RG, Howard EW, Lucroy MD. Effect of wavelength on low-intensity laser irradiation-stimulated cell proliferation in vitro. *Lasers Surg Med* 2005;36:8–12.
- Fonseca AS, Moreira TO, Paixão DL, et al. Effect of laser therapy on DNA damage. *Lasers Surg Med* 2010;42:481–488.
- Hu WP, Wang JJ, Yu CL, Lan CC, Chen GS, Yu HS. Helium-neon laser irradiation stimulates cell proliferation through photostimulatory effects in mitochondria. *J Invest Dermatol* 2007;127:2048–2057.
- Mvula B, Mathope T, Moore TJ, Abrahamse H. The effects of low level laser irradiation on human adipose derived stem cells. *Laser Med Sci* 2008;23:277–282.
- Abrahamse H, Houreld NN, Muller S, Ndlovu L. Fluence and wavelength of low intensity laser irradiation affect activity and proliferation of human adipose derived stem cells. *Medical Technology SA* 2010;24:9–14.
- Chow RT, David MA, Armati PJ. 830 nm laser irradiation induces varicosity formation, reduces mitochondrial membrane potential and blocks fast axonal flow in small and medium diameter rat dorsal root ganglion neurons: implications for the analgesic effects of 830 nm laser. *J Peripher Nerv Syst* 2007;12:28–39.
- Michael R. Hamblin. Mechanisms of low level light therapy, 2008 Available at: <http://www.photobiology.info/Hamblin.html> (Last accessed June 1, 2015).
- Maloney R. Light therapy: low-level laser therapy: treatment through precise application of light. *Bio-Optics World* 2014 Available at: <http://www.bioopticsworld.com/articles/print/volume-7/issue-2/features/light-therapy-low-level-laser-therapy-treatment-through-precise-application-of-light.html> (Last accessed June 1, 2015).
- Tirino V, Desideri V, Paino F, Papaccio G, De Rosa M. Methods for cancer stem cell detection and isolation. *Methods Mol Biol* 2012;879:513–529.
- Salama R, Tang J, Gadgeel SM, Ahmad A, Sarkar FH. Lung cancer stem cells: current progress and future perspectives. *J Stem Cell Res Ther* 2012;S7:007.
- Ren F, Sheng WQ, Du X. CD133: a cancer stem cells marker, is used in colorectal cancers. *World J Gastroenterol* 2013;19:2603–2611.

21. Tirino V, Camerlingo R, Franco R, et al. The role of CD133 in the identification and characterisation of tumour-initiating cells in non-small-cell lung cancer. *Eur J Cardiothorac Surg* 2009;36:446–453.
22. SKUT1 (ATCC® HTB114™) Product sheet. Available at: <https://www.atcc.org/~ps/HTB-114.ashx> (Last accessed April 12, 2014).
23. Manoto SL, Abrahamse H. Effect of a newly synthesized Zn sulfophthalocyanine derivative on cell morphology, viability, proliferation, and cytotoxicity in a human lung cancer cell line (A549). *Lasers Med Sci* 2011;26:523–530.
24. Steynberg T, Visagie MH, Mqoco T, et al. Qualitative assessment of smooth muscle cells propagated on 2d-and 3dpolycaprolactone polymers via scanning electron microscope. *Biomed Res* 2012;23:191–198.
25. Hardt O, Parker LL, Dudley ME, Bosio A. Isolation of tumor cell subpopulations using semi-automated tissue dissociation and magnetic cell separation. *Cancer Res* 2010;70:3357.
26. Eramo A, Lotti F, Sette G, et al. Identification and expansion of the tumorigenic lung cancer stem cell population. *Cell Death Differ* 2008;15:504–514.
27. Promega. CellTiter-Glo® Luminescent Cell Viability Assay, 2012 Available at: <https://worldwide.promega.com/~media/Files/Resources/Protocols/Technical%20Bulletins/0/CellTiter%20Glo%20Luminescent%20Cell%20Viability%20Assay%20Protocol.pdf> (Last accessed July 12, 2014).
28. Rahman M, Deleyrolle L, Vedam-Mai V, Azari H, Abd-El-Barr M, Reynolds BA. The cancer stem cell hypothesis: failures and pitfalls. *Neurosurgery* 2011;68:531–545.
29. Chen YC, Hsu HS, Chen YW, et al. Oct-4 expression maintained cancer stem-like properties in lung cancer-derived CD133-positive cells. *PLoS One* 2008;3:e2637.
30. Takenobu H, Shimoza T, Nakamura T, et al. CD133 Suppresses neuroblastoma cell differentiation via signal pathway modification. *Oncogene* 2011;30:97–105.
31. Caliguri V. The cancer stem cell hypothesis. *BioFiles*, 2008 Available at: <http://www.sigmaaldrich.com/technical-documents/articles/biofiles/the-cancer-stem-cell.html> (Last accessed June 10, 2014).

Address correspondence to:
*Heidi Abrahamse
Laser Research Centre
Faculty of Health Sciences
University of Johannesburg
PO Box 17011
Johannesburg 2028
South Africa*

E-mail: habrahamse@uj.ac.za