

Classes and Theories of Trees Associated with a Class of Linear Orders

VALENTIN GORANKO¹ AND RUAAN KELLERMAN²

¹Department of Informatics and Mathematical Modelling,
Technical University of Denmark

²Department of Mathematics and Applied Mathematics,
University of Pretoria, South Africa

Emails: vfgo@imm.dtu.dk, ruaan.kellerman@up.ac.za

August 27, 2010

Abstract

Given a class of linear order types \mathcal{C} , we identify and study several different classes of trees, naturally associated with \mathcal{C} in terms of how the paths in those trees are related to the order types belonging to \mathcal{C} . We investigate and completely determine the set-theoretic relationships between these classes of trees and between their corresponding first-order theories. We then obtain some general results about the axiomatization of the first-order theories of some of these classes of trees in terms of the first-order theory of the generating class \mathcal{C} , and indicate the problems obstructing such general results for the other classes. These problems arise from the possible existence of non-definable paths in trees, that need not satisfy the first-order theory of \mathcal{C} , so we have started analyzing first-order definable and undefinable paths in trees.

Keywords: *trees, linear orders, first-order theories, definable and non-definable paths*

1 Introduction

Trees occur naturally in many mathematical settings as important partial orders. Every tree is naturally associated with a set of linear orders representing its paths. Conversely, with every set \mathcal{C} of linear order types one can

associate various classes of trees, the paths of which belong, or are related, to \mathcal{C} . Trees are far more complex structures than linear orders, and it is interesting and important to analyze the structural and logical relationships between trees and their sets of paths. While much is known about some important classes of trees and their first-order theories, such as finite trees (see [1]), well-founded trees (see [2]), and finitely branching trees (see [7], [4]), the only systematic study so far, of which we are aware, of classes of trees that naturally arise from a given set of linear order types is [5], in the context of temporal logics. The objective of this paper is to continue and extend that line of research by investigating the variety of classes of trees, the paths in which are naturally associated with a given class of linear orders, and of the first-order theories of these classes. In a subsequent work we apply present results to obtain complete axiomatizations of the first-order theories of some important classes of trees.

The paper is structured as follows: after a preliminary Section 2 we introduce in Section 3 several classes of trees associated in terms of the paths in those trees with a given class of linear orders. Then, in Section 4 we analyze and completely describe the set-theoretic relationships between these classes of trees, whereas in Section 5 we completely describe the relationships between their respective first-order theories. In Section 6 we discuss the problem of axiomatizing these theories given a first-order axiomatization of the generating class of linear orders. In Section 7 we take a closer look at the different types of paths emerging in trees associated with a given class of linear orders and how they are related with the first-order properties of such trees. We end with brief concluding remarks summarizing the paper and discussing the arising open problems and further research directions.

2 Preliminaries

Let \mathfrak{A} be a structure and let $\varphi(x_1, \dots, x_n)$ be a first-order formula. The domain of \mathfrak{A} is denoted as $|\mathfrak{A}|$ or simply as A . Let $c_1, \dots, c_n \in |\mathfrak{A}|$ and put $\bar{x} = (x_1, \dots, x_n)$ and $\bar{c} = (c_1, \dots, c_n)$. Then $\varphi(x_1, \dots, x_n)$ is also written as $\varphi(\bar{x})$. When evaluating the truth of φ in \mathfrak{A} when the elements c_i are substituted for x_i for every i ($1 \leq i \leq n$), we also denote the expression $\mathfrak{A} \models \varphi(c_1/x_1, \dots, c_n/x_n)$ as $\mathfrak{A} \models \varphi(\bar{c}/\bar{x})$. When enriching the signature of \mathfrak{A} with c_1, \dots, c_n as parameters, we also denote $(\mathfrak{A}; c_1, \dots, c_n)$ as $(\mathfrak{A}; \bar{c})$ and $(\mathfrak{A}; \bar{c}) \models \varphi(c_1, \dots, c_n)$ as $(\mathfrak{A}; \bar{c}) \models \varphi(\bar{c})$.

For $\bar{a} = (a_1, \dots, a_k)$ and $\bar{b} = (b_1, \dots, b_n)$, the notation $\bar{a}\bar{c}$ indicates the $(k+1)$ -tuple (a_1, \dots, a_k, c) and $\bar{a}\bar{b}$ indicates the $(k+n)$ -tuple $(a_1, \dots, a_k, b_1, \dots, b_n)$.

The quantifier rank of a formula φ is denoted as $\text{qr}(\varphi)$. Elementary equivalence between structures is denoted by \equiv and n -equivalence (equivalence with respect to all sentences of quantifier rank at most n) by \equiv_n .

2.1 Relativising first-order formulas

Relativisations give a neat method for imposing first-order properties on definable substructures of a structure. The following definition and results are taken from [6, pp. 259-260].

Let \mathfrak{A} be any structure and let $a_1, \dots, a_k \in |A|$. Fix $\bar{x} = (x_1, \dots, x_n)$, $\bar{y} = (y_1, \dots, y_k)$ and $\bar{a} = (a_1, \dots, a_k)$.

Definition 1 ([6]) *Let $\varphi(\bar{x})$ and $\theta(u, \bar{y})$ be any first-order formulas. The relativisation of φ to θ , denoted φ^θ (where $\varphi^\theta = \varphi^\theta(\bar{x}, \bar{y})$), is defined as follows:*

- (i) if φ is atomic then $\varphi^\theta := \varphi$;
- (ii) if $\varphi = \neg\psi$ then $\varphi^\theta := \neg(\psi^\theta)$;
- (iii) if $\varphi = \psi_1 \star \psi_2$ then $\varphi^\theta := \psi_1^\theta \star \psi_2^\theta$, where \star is any of $\vee, \wedge, \rightarrow$ or \leftrightarrow ;
- (iv) if $\varphi = \exists x\psi$ then $\varphi^\theta := \exists x(\theta(x, \bar{y}) \wedge \psi^\theta)$;
- (v) if $\varphi = \forall x\psi$ then $\varphi^\theta := \forall x(\theta(x, \bar{y}) \rightarrow \psi^\theta)$.

Note that if φ is quantifier-free then φ^θ contains the variables y_1, \dots, y_k vacuously, while if φ contains quantifiers then the variables y_1, \dots, y_k will appear explicitly in φ^θ .

Define

$$(\mathfrak{A}; \bar{a})^\theta := \{b \in |A| : (\mathfrak{A}; \bar{a}) \models \theta(b/u, \bar{a})\}.$$

Proposition 2 ([6]) *Let $\varphi(\bar{x})$ and $\theta(u, \bar{y})$ be any first-order formulas. For any $b_1, \dots, b_n \in (\mathfrak{A}; \bar{a})^\theta$ and with $\bar{b} = (b_1, \dots, b_n)$,*

$$\mathfrak{A} \models \varphi^\theta(\bar{b}/\bar{x}, \bar{a}/\bar{y}) \Leftrightarrow (\mathfrak{A}; \bar{a})^\theta \models \varphi(\bar{b}/\bar{x}).$$

Corollary 3 ([6]) *Let \mathfrak{A} be a structure, let σ be a first-order sentence and let $\theta(u, \bar{y})$ be a first-order formula. Then*

$$\mathfrak{A} \models \sigma^\theta(\bar{a}/\bar{y}) \Leftrightarrow (\mathfrak{A}; \bar{a})^\theta \models \sigma.$$

Corollary 4 ([6]) *Let \mathfrak{A} be a structure, let σ be a first-order sentence and let $\theta(u)$ be a first-order formula. Then*

$$\mathfrak{A} \models \sigma^\theta \Leftrightarrow \mathfrak{A}^\theta \models \sigma.$$

2.2 Characteristic formulas

Characteristic formulas give a syntactic formalisation of the Ehrenfeucht-Fraïssé game played on a pair of structures. The following definition and results are borrowed from [2]. An excellent account of characteristic formulas can also be found in [3].

Fix structures \mathfrak{A} and \mathfrak{B} . Let $\bar{a} = (a_1, \dots, a_k)$ and $\bar{b} = (b_1, \dots, b_k)$, where $a_1, \dots, a_k \in |\mathfrak{A}|$ and $b_1, \dots, b_k \in |\mathfrak{B}|$. Put $\bar{x} = (x_1, \dots, x_k)$.

Definition 5 ([2]) For $n \in \mathbb{N}$ we define the formula $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n$ (with $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n = \llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n(\bar{x})$) inductively as follows:

- (i) $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^0 := \bigwedge \{ \varphi(\bar{x}) : \varphi \text{ an atomic or negated atomic formula with } \mathfrak{A} \models \varphi(\bar{a}/\bar{x}) \};$
- (ii) $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^{m+1} := \bigwedge_{a_{k+1} \in |\mathfrak{A}|} \exists x_{k+1} \llbracket (\mathfrak{A}; \bar{a}a_{k+1}) \rrbracket^m \wedge \forall x_{k+1} \bigvee_{a_{k+1} \in |\mathfrak{A}|} \llbracket (\mathfrak{A}; \bar{a}a_{k+1}) \rrbracket^m.$

The formula $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n$ is known as the n -characteristic of \bar{a} in \mathfrak{A} .

The formula $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n$ describes any (the first conjunct in clause (ii) of Definition 5) and all (the second conjunct in clause (ii) of Definition 5) ways to relate n -tuples of elements from \mathfrak{A} to the elements in \bar{a} .

Lemma 6 ([2])

- (i) $\mathfrak{A} \models \llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n(\bar{a}/\bar{x});$
- (ii) the formula $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n$ has quantifier rank n .

Theorem 7 ([2]) For $n \in \mathbb{N}$ the following conditions are equivalent:

- (i) $(\mathfrak{A}; \bar{a}) \equiv_n (\mathfrak{B}; \bar{b});$
- (ii) $\mathfrak{B} \models \llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n(\bar{b}/\bar{x});$
- (iii) the formulas $\llbracket (\mathfrak{A}; \bar{a}) \rrbracket^n$ and $\llbracket (\mathfrak{B}; \bar{b}) \rrbracket^n$ are equivalent.

Corollary 8 Let $n \in \mathbb{N}$. The following conditions are equivalent:

- (i) $\mathfrak{A} \equiv_n \mathfrak{B};$
- (ii) $\mathfrak{B} \models \llbracket \mathfrak{A} \rrbracket^n;$
- (iii) the formulas $\llbracket \mathfrak{A} \rrbracket^n$ and $\llbracket \mathfrak{B} \rrbracket^n$ are equivalent.

Hence the n -characteristics of empty tuples are canonical objects associated with classes of structures which are n -equivalent.

When working with a finite signature, there will be only finitely many n -characteristics of k -tuples.

Theorem 9 ([2]) *Let $\{\mathfrak{A}_i\}_{i \in I}$ be a class of structures over the same finite signature. Let $n, k \in \mathbb{N}$ and let \bar{a}_i be a k -tuple from \mathfrak{A}_i for every $i \in I$. The set of pairwise non-equivalent formulas $\llbracket (\mathfrak{A}_i; \bar{a}_i) \rrbracket^n$ ($i \in I$) is finite.*

2.3 Trees

A *tree* is a partially ordered set $(T; <)$ satisfying the following two properties:

- (i) for every $x \in T$ the set $\{y \in T : y < x\}$ is totally ordered, and
- (ii) for every $x, y \in T$, there exists $z \in T$ such that $z \leq x, y$.

Thus the class of trees can be defined in the language containing the binary relation $<$ together with the usual equality relation $=$, using the first-order theory consisting of the following sentences:

$$A_1: \forall x (\neg x < x);$$

$$A_2: \forall x \forall y \forall z (x < y \wedge y < z \rightarrow x < z);$$

$$A_3: \forall x \forall y \forall z (y < x \wedge z < x \rightarrow (x < y \vee x = y \vee y < x));$$

$$A_4: \forall x \forall y \exists z ((z < x \vee z = x) \wedge (z < y \vee z = y)).$$

We will denote the first-order theory axiomatized with the four sentences above by **Tree**.

When the context allows we will simply write T for $(T; <)$. Also, we write $x \leq y$ as a shorthand for $x < y \vee x = y$. The elements of a tree are called *nodes*. If $a < b$ then a is called the *predecessor* of b and b is called a *successor* of a ; if there is no c with $a < c < b$ then a is called the *immediate predecessor* of b and b is called an *immediate successor* of a . A tree is called *binary* if every node in it has precisely two immediate successors. A $<$ -maximal node is called a *leaf*. The set of leaves can be defined using the first-order formula $\text{leaf}(x) := \forall y (x \leq y \rightarrow x = y)$.

A maximal totally ordered set of nodes is called a *path*. Using Zorn's Lemma, it is easy to see that every totally ordered subset of a tree is contained in a path. For any order type α , a path A in a tree is called an α -*path* if A is isomorphic with α . The binary tree in which every path is an α -path will be denoted B_α . As usual, we denote by ω the order type of the positive

integers, by ω^* the order type of the negative integers, and by ζ the order type of all integers.

A tree T is called *well-founded* when every non-empty set of nodes from T contains a minimal node.

A tree is called *downwards discrete* when every non-root node in T has an immediate predecessor. T is called *weakly upwards discrete* when every non-leaf node has an immediate successor, and *upwards discrete* when, for every path X in T , every non-leaf node in X has an immediate successor belonging to X . T is called *weakly discrete* when it is both downwards discrete as well as weakly upwards discrete, and *discrete* when it is both downwards discrete and upwards discrete. Note that all these properties are first-order definable.

A discrete tree is called *finitely branching* when every node in that tree has finitely many (possibly none) immediate successors.¹

Given a tree T and $a \in T$, define

$$\begin{aligned} a_{>} &:= \{x \in T : x < a\}, \\ a_{\geq} &:= \{x \in T : x \leq a\}, \\ a_{<} &:= \{x \in T : a < x\}, \\ a_{\leq} &:= \{x \in T : a \leq x\}. \end{aligned}$$

The sets $a_{>}$, a_{\geq} , $a_{<}$ and a_{\leq} will also be treated as substructures of T . The sets $a_{>}$ and a_{\geq} are always linear orders, while the set a_{\leq} is always a tree – the *generated subtree of T rooted at a* .

For $k \in \mathbb{N}$ and $\varphi(x, \bar{z})$ any formula with $\bar{z} = (z_1, \dots, z_k)$, define the formula

$$\begin{aligned} \pi_\varphi(\bar{z}) := & \exists x \varphi(x, \bar{z}) \wedge \forall x \forall y (\varphi(x, \bar{z}) \wedge \varphi(y, \bar{z}) \rightarrow (x < y \vee x = y \vee y < x)) \wedge \\ & \forall x \forall y (x < y \wedge \varphi(y, \bar{z}) \rightarrow \varphi(x, \bar{z})) \wedge \neg \exists x \forall y (\varphi(y, \bar{z}) \rightarrow y < x). \end{aligned}$$

Note that, if φ has quantifier rank n , then π_φ has quantifier rank $n + 2$.

The formula π_φ formalizes the claim that the formula φ defines a path. Verifying that is straightforward, thus obtaining the following.

Proposition 10 *Let T be a tree with $c_1, \dots, c_k \in T$ and put $\bar{c} = (c_1, \dots, c_k)$. The formula $\varphi(x, \bar{c})$ defines a path in $(T; \bar{c})$ if and only if $T \models \pi_\varphi(\bar{c}/\bar{z})$.*

By the proposition above, if $T_1 \equiv T_2$ then a formula φ defines a path in T_1 iff it defines a path in T_2 .

¹This definition of finite branching is not the most general, nor necessarily the most intuitively acceptable, but it is sufficient for the purposes of this paper. For further details and discussion on finite branching see [7] and [4].

A substructure S of a tree T which is itself a tree is called a *subtree* of T . S is called *upwards closed* in T when it satisfies the property that for all $x, y \in T$, whenever $x \in S$ and $x < y$ then $y \in S$.

Proposition 11 ([4]) *Let $T_1 = (T_1; <_{T_1})$ be a tree and let $\{(S_i; <_{S_i}) : i \in I\}$ be a pairwise disjoint set of upwards closed subtrees of T_1 , where the order $<_{S_i}$ is the order $<_{T_1}$ restricted to the domain S_i . For every $i \in I$, let $S_{f(i)} = (S_{f(i)}; <_{S_{f(i)}})$ be a tree with $S_i \equiv_n S_{f(i)}$. Let T_2 be the tree obtained from T_1 by replacing every subtree S_i with the tree $S_{f(i)}$. Formally we define $T_2 = (T_2; <_{T_2})$ as follows:*

- $|T_2| := (T_1 \setminus \bigcup_{i \in I} S_i) \cup \bigcup_{i \in I} S_{f(i)}$, and
- for $x, y \in T_2$, $x <_{T_2} y$ if and only if one of the following conditions are satisfied:
 - (i) $x, y \in T_1 \setminus \bigcup_{i \in I} S_i$ and $x <_{T_1} y$, or
 - (ii) $x, y \in S_{f(i)}$ for some i and $x <_{S_{f(i)}} y$, or
 - (iii) $x \in T_1 \setminus S_i$ and $y \in S_{f(i)}$ for some i , and $x <_{T_1} z$ for some $z \in S_i$.

Then $T_1 \equiv_n T_2$.

3 Classes of trees associated with a given class of linear orders

Here we define several classes of trees naturally associated, from the viewpoint of first-order logic, with a given set of linear order types \mathcal{C} . We will call these classes of trees \mathcal{C} -classes.

Definition 12 *A tree T is called a:*

- (i) \mathcal{C} -tree when every path in T is an α -path for some $\alpha \in \mathcal{C}$;
- (ii) uniformly \mathcal{C} -like tree (U- \mathcal{C} -like tree) if $T \equiv S$ for some \mathcal{C} -tree S ;
- (iii) \mathcal{C} -like tree if, for every $n \in \mathbb{N}$, there is a \mathcal{C} -tree S such that $T \equiv_n S$;
- (iv) pathwise uniformly \mathcal{C} -like tree (PU- \mathcal{C} -like tree) if, for every path X in T , there exists $\alpha \in \mathcal{C}$ such that $X \equiv \alpha$;
- (v) pathwise \mathcal{C} -like tree (P- \mathcal{C} -like tree) if, for every path X in T and for every $n \in \mathbb{N}$, there exists $\alpha \in \mathcal{C}$ such that $X \equiv_n \alpha$;

- (vi) definably \mathcal{C} -tree (D- \mathcal{C} -tree) if every parametrically definable path X in T is an α -path for some $\alpha \in \mathcal{C}$ (dependent on X);
- (vii) definably uniformly \mathcal{C} -like tree (DU- \mathcal{C} -like tree) if, for every parametrically definable path X in T , there exists $\alpha \in \mathcal{C}$ such that $X \equiv \alpha$;
- (viii) definably \mathcal{C} -like tree (D- \mathcal{C} -like tree) if, for every parametrically definable path X in T and for every $n \in \mathbb{N}$, there exists $\alpha \in \mathcal{C}$ such that $X \equiv_n \alpha$. Equivalently (since the language of trees has finite signature), if every parametrically definable path in T is a model of the first-order theory of \mathcal{C} .

If $\mathcal{C} = \{\alpha\}$ then T is simply called an α -tree, a uniformly α -like tree, etc.

We follow with a few examples that will be used further, in the proof of Theorem 19.

Example 13 The tree $B_{\omega+1}$ is uncountable. Let T_1 be any countable elementary substructure of $B_{\omega+1}$ and let $\mathcal{C}_1 = \{\omega + \mathbf{1}\}$. T_1 is a binary tree not containing any finite paths, so that every path in T_1 is either an ω -path or an $(\omega + \mathbf{1})$ -path. Moreover, T_1 does actually contain both ω -paths and $(\omega + \mathbf{1})$ -paths (and the former are not even 2-equivalent to $\omega + \mathbf{1}$). Thus, T_1 can be seen as the result of removing an uncountable set of leaves from $B_{\omega+1}$.

Suppose now, that $\varphi(x, \bar{c})$ defines a path A in $(T_1; \bar{c})$ for some tuple \bar{c} of nodes from T_1 . Then $(T_1, \bar{c}) \models \pi_\varphi(\bar{c})$, so $(B_{\omega+1}, \bar{c}) \models \pi_\varphi(\bar{c})$, hence $\varphi(x, \bar{c})$ defines a path in $(B_{\omega+1}; \bar{c})$. Now, since every path in $B_{\omega+1}$ is an $(\omega + \mathbf{1})$ -path then we get that $(B_{\omega+1}, \bar{c}) \models \exists x (\text{leaf}(x) \wedge \varphi(x, \bar{c}))$ and so $(T_1, \bar{c}) \models \exists x (\text{leaf}(x) \wedge \varphi(x, \bar{c}))$. Hence A will contain a leaf. Thus every parametrically definable path in T_1 will contain a leaf, and since every path in T_1 containing a leaf is parametrically definable (using that leaf as parameter) it follows that the parametrically definable paths in T_1 are precisely its $(\omega + \mathbf{1})$ -paths.

Thus, T_1 is a uniformly $(\omega + \mathbf{1})$ -like tree, as well as a definably $(\omega + \mathbf{1})$ -tree, but neither an $(\omega + \mathbf{1})$ -tree, nor a pathwise $(\omega + \mathbf{1})$ -like tree.

Example 14 Let T_2 be the tree indicated in Figure 1 and let $\mathcal{C}_2 = \{\omega\}$. Each of the two paths in T_2 is parametrically definable and elementarily equivalent with ω . In any ω -like tree, every parametrically definable set contains a minimal node. The set of nodes in T_2 defined by the formula

$$\varphi(x) = \forall y \forall z (x < y \wedge x < z \rightarrow (y < z \vee y = z \vee z < y))$$

contains no minimal node. Thus, T_2 is a definably uniformly ω -like tree, but not ω -like.



Figure 1: The trees T_2 and T_3 described in Example 14 and Example 15.

Let σ_1 be the sentence

$$\sigma_1 := \exists u \varphi(u) \rightarrow \exists u (\varphi(u) \wedge \forall w (w < u \rightarrow \neg \varphi(w))),$$

where $\varphi(x)$ is defined as above. This sentence will be used further.

Example 15 Let T_3 be the tree indicated in Figure 1 and let $\mathcal{C}_3 = \{\mathbf{n} : n \in \mathbb{N}\}$. Both paths in T_3 are parametrically definable. It is known (e.g, [6]) that for every m there exists some sufficiently large n such that $\omega + \omega^* \equiv_m \mathbf{n}$. However, $\omega + \omega^* \not\equiv \mathbf{n}$ for every n . In any $DU\text{-}\mathcal{C}_3$ -like tree, the set defined by the formula $\varphi(x)$ from Example 14 will contain a minimal node. However, the subset of T_3 defined by $\varphi(x)$ does not contain a minimal node. Thus, T_3 is a definably \mathcal{C}_3 -like tree, even a pathwise \mathcal{C}_3 -like tree, but neither a \mathcal{C}_3 -like tree nor a definably uniformly \mathcal{C}_3 -like tree.

Example 16 Let T_4 be the linear order $\omega + \omega^*$ and let $\mathcal{C}_4 = \{\mathbf{n} : n \in \mathbb{N}\}$. As noted above, there exists, for every m , some sufficiently large n such that $\omega + \omega^* \equiv_m \mathbf{n}$, but $\omega + \omega^* \not\equiv \mathbf{n}$ for every n .

Example 17 Let T_5 be the tree B_ω and take $\mathcal{C}_5 = \{\omega + \mathbf{1}\}$. Note that B_ω contains no parametrically definable paths. Let σ_2 be the sentence

$$\sigma_2 := \forall x \exists y (x \leq y \wedge \text{leaf}(y)).$$

This sentence will be used further.

Example 18 Let T_6 be the linear order $\omega + \zeta$ and take $\mathcal{C}_6 = \{\omega\}$. It is known that $\omega \equiv \omega + \zeta$.




Figure 2: Relationships between the \mathcal{C} -classes of trees (see Theorem 19). Inclusions $X \subseteq Y$ are denoted as $X \rightarrow Y$. Non-inclusions are indicated by specifying a counterexample drawn from Examples 13 - 18 or, when obtained through completion of diagrams, by \times . There are no downwards directed inclusions between classes separated by more than one level.

4 Relationships between the \mathcal{C} -classes of trees

Theorem 19 *Let \mathcal{C} be a class of linear orders. The set-theoretical inclusions and non-inclusions that hold between the various \mathcal{C} -classes of trees are presented in Figure 2.*

Proof

To begin with the inclusions, we will show that the class of \mathcal{C} -like trees is contained in the class of D- \mathcal{C} -like trees. The argument to show that the class of U- \mathcal{C} -like trees is contained in the class of DU- \mathcal{C} -like trees is similar. The remaining inclusions are quite easy to verify.

Let T be a \mathcal{C} -like tree and let A be a path in T defined in $(T; \bar{c})$ by the formula $\varphi(x, \bar{c})$ for some tuple of nodes \bar{c} from T . Suppose that A has n -characteristic τ . Then $T \models \pi_\varphi(\bar{c}/\bar{z})$ and $T \models \tau^\varphi(\bar{c}/\bar{z})$ so that $T \models \exists \bar{z} (\pi_\varphi(\bar{z}) \wedge \tau^\varphi(\bar{z}))$. Since T is \mathcal{C} -like then there exists a \mathcal{C} -tree S for

which $S \models \exists \bar{z} (\pi_\varphi(\bar{z}) \wedge \tau^\varphi(\bar{z}))$. Thus $\varphi(x, \bar{d})$ defines a path B in $(S; \bar{d})$ for some tuple \bar{d} from S , and $B \models \tau$. But B is isomorphic with some linear order C in \mathcal{C} and so $A \equiv_n C$. It follows that T is a $D\text{-}\mathcal{C}$ -like tree.

As an example of a non-inclusion demonstrated by a counterexample, we show that the class of $P\text{-}\mathcal{C}$ -like trees is not always included in the class of \mathcal{C} -like trees. Note, that the tree T_2 from Example 14 is a $P\text{-}\mathcal{C}_2$ -like tree, but not a \mathcal{C}_2 -like tree. This is because every \mathcal{C}_2 -like tree satisfies the sentence σ_1 as defined in Example 14, while T_2 does not. Hence, the class of $P\text{-}\mathcal{C}_2$ -like trees is not contained in the class of \mathcal{C}_2 -like trees.

As an example of a non-inclusion obtained through transitive completion in Figure 2, consider the claim that the class of $P\text{-}\mathcal{C}$ -like trees is not generally a subclass of the class of $PU\text{-}\mathcal{C}$ -like trees. If, to the contrary, the class of $P\text{-}\mathcal{C}$ -like trees were a subclass of the class of $PU\text{-}\mathcal{C}$ -like trees for all classes of linear orders \mathcal{C} , then since the class of $PU\text{-}\mathcal{C}$ -like trees is also a subclass of the class of $DU\text{-}\mathcal{C}$ -like trees for all classes \mathcal{C} , we would get that the class of $P\text{-}\mathcal{C}$ -like trees is a subclass of the class of $DU\text{-}\mathcal{C}$ -like trees for all classes \mathcal{C} . But this contradicts the fact that the tree T_4 from Example 16 is a $P\text{-}\mathcal{C}_4$ -like tree, with \mathcal{C}_4 as defined in Example 16, but T_4 is not a $DU\text{-}\mathcal{C}_4$ -like tree. This establishes the non-inclusion.

The remaining non-inclusions are easily verified. –

The verification of the claims in the following proposition is routine.

Proposition 20 *When the class \mathcal{C} consists of a single linear order, the following additional inclusions hold:*

- (i) *the class of $P\text{-}\mathcal{C}$ -like trees \subseteq the class of $PU\text{-}\mathcal{C}$ -like trees,*
- (ii) *the class of $D\text{-}\mathcal{C}$ -like trees \subseteq the class of $DU\text{-}\mathcal{C}$ -like trees.*

Consequently:

- (iii) *the class of \mathcal{C} -like trees \subseteq the class of $DU\text{-}\mathcal{C}$ -like trees,*
- (iv) *the class of $P\text{-}\mathcal{C}$ -like trees \subseteq the class of $DU\text{-}\mathcal{C}$ -like trees.*

5 Relationships between the first-order theories of \mathcal{C} -classes of trees

Theorem 21 *Let \mathcal{C} be a class of linear orders. The set-theoretical inclusions and non-inclusions that hold between the first-order theories of the various \mathcal{C} -classes of trees are presented in Figure 3.*




Figure 3: Relationships between the first-order theories of the \mathcal{C} -classes of trees (see Theorem 21). Inclusions $X \subseteq Y$ are denoted as $X \rightarrow Y$. Non-inclusions are indicated by specifying a counterexample drawn from Examples 13 - 18 or, when obtained through completion, by the symbol \times . Except for the obvious cases, there are no upwards directed inclusions between the theories of classes separated by more than one level.

Proof Let $\sigma \in \text{TH}(\text{P-}\mathcal{C}\text{-like trees})$ say with $\text{qr}(\sigma) = n$. Let T be a \mathcal{C} -like tree. Then $T \equiv_n T_0$ for some \mathcal{C} -tree T_0 . But T_0 is also a P- \mathcal{C} -like tree hence $T_0 \models \sigma$ and so $T \models \sigma$. It follows that $\sigma \in \text{TH}(\mathcal{C}\text{-like trees})$ and so $\text{TH}(\text{P-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\mathcal{C}\text{-like trees})$.

The following inclusions can be proven using a similar argument:

- $\text{TH}(\text{DU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{D-}\mathcal{C}\text{-trees}) \subseteq \text{TH}(\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{PU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{U-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{PU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{U-}\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{P-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{U-}\mathcal{C}\text{-like trees})$,
- $\text{TH}(\text{D-}\mathcal{C}\text{-trees}) \subseteq \text{TH}(\text{U-}\mathcal{C}\text{-like trees})$.

The inclusion $\text{TH}(\mathcal{C}\text{-trees}) \subseteq \text{TH}(\text{U-}\mathcal{C}\text{-like trees})$ is immediate.

The remaining inclusions follow from Theorem 19 and the accompanying diagram in Figure 2.

We briefly discuss the non-inclusions shown in Figure 3. Consider for example the non-inclusion $\text{TH}(\text{DU-}\mathcal{C}_3\text{-like trees}) \not\subseteq \text{TH}(\text{P-}\mathcal{C}_3\text{-like trees})$. To see this, let T be a definably uniformly \mathcal{C}_3 -like tree and suppose that $T \models \varphi(a/x)$ for some $a \in T$, where $\varphi(x)$ is as in Example 14. Then a belongs to a parametrically definable (being 'singular', see next section) path A , with $A \equiv \mathbf{n}$ for some $n \in \mathbb{N}$. Hence A will be finite. It follows that $T \models \sigma_1$ (where σ_1 is as in Example 14) and so $\sigma_1 \in \text{TH}(\text{DU-}\mathcal{C}_3\text{-trees})$. However, T_3 is a P- \mathcal{C}_3 -like tree with $T_3 \not\models \sigma_1$, so that $\sigma_1 \notin \text{TH}(\text{P-}\mathcal{C}_3\text{-like trees})$. This serves as a counterexample to establish the non-inclusion $\text{TH}(\text{DU-}\mathcal{C}_3\text{-like trees}) \not\subseteq \text{TH}(\text{P-}\mathcal{C}_3\text{-like trees})$.

It can be shown, using an argument similar to the one in the preceding paragraph, that $\text{TH}(\text{D-}\mathcal{C}_2\text{-trees}) \not\subseteq \text{TH}(\text{PU-}\mathcal{C}_2\text{-like trees})$.

The non-inclusions which use the class \mathcal{C}_5 and the sentence σ_2 from Example 17 as counterexample are easily verified.

Finally, the non-inclusions obtained through completion are trivial. For example, $\text{TH}(\mathcal{C}\text{-trees})$ is not generally a subtheory of $\text{TH}(\text{D-}\mathcal{C}\text{-trees})$, for if it were, then using the fact that $\text{TH}(\text{PU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\mathcal{C}\text{-trees})$ for all classes \mathcal{C} , this would give $\text{TH}(\text{PU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{D-}\mathcal{C}\text{-trees})$ for all classes \mathcal{C} , contradicting the fact that $\text{TH}(\text{PU-}\mathcal{C}_5\text{-like trees}) \not\subseteq \text{TH}(\text{D-}\mathcal{C}_5\text{-trees})$.

Likewise the theory $\text{TH}(\mathcal{C}\text{-like trees})$ is not generally a subtheory of the theory $\text{TH}(\text{D-}\mathcal{C}\text{-trees})$, for if it were, then the theories $\text{TH}(\text{D-}\mathcal{C}\text{-trees})$, $\text{TH}(\mathcal{C}\text{-like trees})$ and $\text{TH}(\text{U-}\mathcal{C}\text{-like trees})$ would coincide for all classes \mathcal{C} . But this would contradict the fact that $\text{TH}(\mathcal{C}\text{-trees}) \subseteq \text{TH}(\text{U-}\mathcal{C}\text{-like trees})$ for all classes \mathcal{C} , while there exist classes \mathcal{C} for which $\text{TH}(\mathcal{C}\text{-trees}) \not\subseteq \text{TH}(\text{D-}\mathcal{C}\text{-trees})$.

The remaining non-inclusions can be proved using similar reasoning. \dashv

Proposition 22 *If \mathcal{C} consists of a single linear order, the following additional inclusions hold between the first-order theories of \mathcal{C} -classes of trees:*

- (i) $\text{TH}(\text{DU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{D-}\mathcal{C}\text{-like trees})$,
- (ii) $\text{TH}(\text{DU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{P-}\mathcal{C}\text{-like trees})$,
- (iii) $\text{TH}(\text{PU-}\mathcal{C}\text{-like trees}) \subseteq \text{TH}(\text{P-}\mathcal{C}\text{-like trees})$.

The other non-inclusions remain the same.

6 Axiomatizing the first-order theories of the \mathcal{C} -classes of trees

As shown in the previous section, every class of linear orders generates eight naturally associated classes of trees, which in general have up to six different first-order theories. In this section we discuss the following problem: given a class of linear orders \mathcal{C} , can we determine the first-order theories of the respective \mathcal{C} -classes of trees in terms of the first-order theory of \mathcal{C} ? In principle, this is a rather difficult problem, because \mathcal{C} may consist of a quite arbitrary collection of linear order types. Here we will obtain some general results, for some of the associated classes of trees; more specific results will be included in a follow-up work.

A natural family of axioms for the class of trees constructed from \mathcal{C} arises as follows: every parametrically definable path in every such tree must satisfy all axioms of the theory of \mathcal{C} . To be more precise, for any theory of linear orders Σ , let De_Σ denote the scheme consisting of all sentences

$$\forall \bar{z} (\pi_\varphi(\bar{z}) \rightarrow \sigma^\varphi(\bar{z}))$$

for every formula $\varphi(x, \bar{z})$ (including formulas $\varphi(x)$ for which the tuple \bar{z} is empty) and for every sentence $\sigma \in \Sigma$. If $\Sigma = \{\sigma\}$ then De_Σ is written simply as De_σ . The scheme De_Σ states that every parametrically definable path satisfies the theory Σ .

This axiom scheme axiomatizes the class of definably \mathcal{C} -like trees:

Proposition 23 *Let \mathcal{C} be a class of linear order types axiomatized by the theory Σ . The class of definably \mathcal{C} -like trees is precisely the class of models of the theory $\text{Tree} \cup \text{De}_\Sigma$.*

Proof Let T be a definably \mathcal{C} -like tree. It is immediate that T satisfies **Tree**. Let $\varphi(x, \bar{z})$ be a formula with $\bar{z} = (z_1, \dots, z_k)$ (\bar{z} may be empty), let $\bar{c} = (c_1, \dots, c_k)$ be a tuple of nodes from T , and let $T \models \pi_\varphi(\bar{c}/\bar{z})$. Then there is a path A defined in $(T; \bar{c})$ by $\varphi(x, \bar{c})$. But $A \models \sigma$ for every $\sigma \in \Sigma$ and $A = (T; \bar{c})^\varphi$, so $T \models \sigma^\varphi(\bar{c}/\bar{z})$ for every $\sigma \in \Sigma$. Thus, $T \models \text{De}_\Sigma$.

Conversely, let T be a structure such that $T \models \text{Tree} \cup \text{De}_\Sigma$. Then T is a tree. Let \bar{c} be a (possibly empty) k -tuple of nodes in T , let $\varphi(x, \bar{z})$ be a formula with $\bar{z} = (z_1, \dots, z_k)$, and let A be a path definable in $(T; \bar{c})$ using the formula $\varphi(x, \bar{c})$. Then $T \models \pi_\varphi(\bar{c}/\bar{z})$ hence $T \models \sigma^\varphi(\bar{c}/\bar{z})$ for every $\sigma \in \Sigma$. But $A = (T; \bar{c})^\varphi$ so $A \models \sigma$ for every $\sigma \in \Sigma$. Hence A is a model of $\text{TH}(\mathcal{C})$. Thus, T is a definably \mathcal{C} -like tree. \dashv

When \mathcal{C} is an axiomatizable class of linear order types, the classes of definably \mathcal{C} -like trees and definably \mathcal{C} -trees coincide, hence the following holds.

Corollary 24 *Let \mathcal{C} be an axiomatizable class of linear order types, axiomatized by the theory Σ . Then the class of definably \mathcal{C} -trees is precisely the class of models of the theory $\text{Tree} \cup \text{De}_\Sigma$.*

Likewise, if \mathcal{C} consists of finitely many linear order types, then the classes of definably \mathcal{C} -like trees and definably uniformly \mathcal{C} -like trees coincide, hence the following holds.

Corollary 25 *Let \mathcal{C} be a finite class of linear order types, axiomatized by the theory Σ . Then the class of definably uniformly \mathcal{C} -like trees is precisely the class of models of the theory $\text{Tree} \cup \text{De}_\Sigma$.*

As is well known (see e.g. [6]), for every ordinal α with $\alpha < \omega^\omega$, there exists a first-order sentence Φ_α which axiomatizes the first-order theory of α , and $\Phi_\alpha \equiv \Phi_\beta$ if and only if $\alpha = \beta$. Hence we have the following corollary.

Corollary 26 *Let α be an ordinal with $\alpha < \omega^\omega$. The class of definably (uniformly) α -like trees is precisely the class of models of the theory $\text{Tree} \cup \text{De}_{\Phi_\alpha}$.*

Now, what about the other \mathcal{C} -classes of trees, characterized in terms of the entire structures, or the sets of *all* paths in them, not only the parametrically definable ones? To get a grip on the axiomatizations of such classes one has to analyze the *non-definable* paths emerging in them. We will consider that question in the next section, where we will take a closer look at first-order parametrically definable and undefinable paths in trees.

7 First-order parametrically definable and undefinable paths in trees: zooming in

7.1 On parametrically definable paths

For α an ordinal and T a well-founded tree, a node a in T is said to have *level* α when α is the order type of the set $a_{>}$. The supremum of the set of all levels of nodes in T is called the *height* of T . The set of nodes having level α can be defined using the formula

$$\text{level}_\alpha(y) := \Phi_\alpha^{\theta(x,y)}(y),$$

where $\theta(x, y) := x \leq y$.

The next result shows that in well-founded trees T of height less than ω^ω , the ability of nodes to define subsets (in particular, paths) of T , improves with the level of those nodes.

Proposition 27 *Let T be a well-founded tree of height less than ω^ω . Let $\bar{c} = (c_1, \dots, c_k)$ be a tuple of nodes from T and let $\varphi(x, \bar{z})$ be a formula, with $\bar{z} = (z_1, \dots, z_k)$, such that $\varphi(x, \bar{c})$ defines the set A in $(T; \bar{c})$. For every i ($1 \leq i \leq k$), let $d_i \in T$ with $c_i \leq d_i$. Then there is a formula $\psi(x, \bar{z})$ such that $\psi(x, \bar{d})$ defines A in $(T; \bar{d})$, where $\bar{d} = (d_1, \dots, d_k)$.*

Proof For every i , suppose c_i has level α_i . Then c_i can be defined in $(T; \bar{d})$ using the formula $\gamma_i(y, d_i) := y \leq d_i \wedge \text{level}_{\alpha_i}(y)$. Hence take

$$\psi(x, z_1, \dots, z_k) := \forall y_1 \dots \forall y_k \left(\bigwedge_{i=1}^k \gamma_i(y_i, z_i) \rightarrow \varphi(x, y_1, \dots, y_k) \right).$$

†

In particular, in well-founded trees of height less than ω^ω for which every non-leaf node has a successor which is a leaf, every parametrically definable set of nodes can be defined using leaves for parameters.

Proposition 28 *Let T be a finitely branching tree in which every node has finite level and let A be a path in T . Suppose A is definable in $(T; \bar{c})$ for some tuple of nodes $\bar{c} = (c_1, \dots, c_k)$ from T . Then there exists $d \in A$ such that A is definable in $(T; d)$.*

Proof We first show that the parameter c_k can be replaced with a parameter d_k from A itself. Let $\varphi(x, \bar{z})$ be a formula (with $\bar{z} = (z_1, \dots, z_k)$) such that

$\varphi(x, \bar{c})$ defines A in $(T; \bar{c})$. Suppose φ has quantifier rank n and that c_k has level l . Let

$$B := \left\{ b \in T : T \models \llbracket (T; \bar{c}) \rrbracket^{n+2}(c_1/x_1, \dots, c_{k-1}/x_{k-1}, b/x_k) \text{ and } b \text{ has level } l \right\}.$$

Since $c_k \in B$ we have that B is non-empty. From the fact that T is finitely branching and that every node in T has finite level it follows that B is finite. B can be defined in $(T; c_1, \dots, c_{k-1})$ using the formula $\xi(x, c_1, \dots, c_{k-1})$, where

$$\xi(x, z_1, \dots, z_{k-1}) := \llbracket (T; \bar{c}) \rrbracket^{n+2}(z_1, \dots, z_{k-1}, x) \wedge \text{level}_l(x).$$

Since $(T; \bar{c}) \models \pi_\varphi(\bar{c})$ and $(T; c_1, \dots, c_k) \equiv_{n+2} (T; c_1, \dots, c_{k-1}, b)$ for every $b \in B$ then $\varphi(x, c_1, \dots, c_{k-1}, b)$ defines a path in $(T; c_1, \dots, c_{k-1}, b)$ for every $b \in B$. Hence the formula $\zeta(x, c_1, \dots, c_{k-1})$, where

$$\zeta(x, z_1, \dots, z_{k-1}) := \exists y (\xi(y, z_1, \dots, z_{k-1}) \wedge \varphi(x, z_1, \dots, z_{k-1}, y)),$$

defines in $(T; c_1, \dots, c_{k-1})$ a subtree T_0 of T containing only finitely many paths, amongst which is A .

Now, choose any $d_k \in A$ such that d_k does not belong to any path in T_0 except for A . Then A can be defined in $(T; c_1, \dots, c_{k-1}, d_k)$ using the formula $\chi(x, c_1, \dots, c_{k-1}, d_k)$, where

$$\chi(x, z_1, \dots, z_k) := \zeta(x, z_1, \dots, z_{k-1}) \wedge (x < z_k \vee x = z_k \vee z_k < x).$$

Hence we have succeeded in replacing the parameter c_k with a parameter d_k from A .

Repeating this procedure for the parameters c_{k-1}, \dots, c_1 , we eventually obtain nodes $d_1, \dots, d_k \in A$ and a formula $\chi'(x, z_1, \dots, z_k)$ such that $\chi'(x, d_1, \dots, d_k)$ defines A in $(T; d_1, \dots, d_k)$. Suppose without loss of generality that $d_i \leq d_1$ for every $i \geq 2$ and that the level of d_i is l_i . Then d_i can be defined in $(T; d_1)$ using the formula $x \leq d_1 \wedge \text{level}_{l_i}(x)$. It follows that A can be defined in $(T; d_1)$ using the formula $\psi(x, d_1)$, where

$$\psi(x, z) := \forall z_2 \dots \forall z_k \left(\bigwedge_{i=2}^k (z_i \leq z \wedge \text{level}_{l_i}(z_i)) \rightarrow \chi'(x, z, z_2, \dots, z_k) \right).$$

Now, take $d = d_1$. +

7.2 Singular and emergent paths

It is generally not possible to formalise in the first-order language of trees with equality the requirement that all paths in a given tree satisfy the first-order theory of some class \mathcal{C} of linear orders, and subsequently to axiomatize the first-order theory of the class of \mathcal{C} -trees. This is because not all paths in a tree need be parametrically definable, so we need to analyze deeper the nature of non-parametrically definable paths in trees, too. The behaviour and structure of such paths can be better understood by considering two types of paths: singular and emergent.

Definition 29 *Let T be a tree and let A be a path in T . A is called singular if there exists $a \in A$ such that a_{\leq} is total. Otherwise the path A is called emergent. If \mathcal{B} is a set of paths from T with $A \notin \mathcal{B}$ and with $A \subseteq \bigcup \mathcal{B}$ then A is said to emerge from \mathcal{B} .*

For a more detailed analysis of singular and emergent paths, the reader is referred to [5].

Example 30 *Let T be the tree obtained by taking the linear order $A := \omega$ and at each point in A , we adjoin a copy of ω (see Figure 4). Thus every path in T will be isomorphic with ω . The path A is an emergent path, while every other path in T is singular.*




Figure 4: Singular and emergent paths (see Example 30).

Clearly, every path containing a greatest node is singular.

Every singular path is parametrically definable. Indeed, let A be a singular path in a tree T and let $a \in A$ such that a_{\leq} is total. Then A can be defined in $(T; a)$ using the formula

$$x < a \vee x = a \vee a < x.$$

It hence follows that, given any tree T , the set of paths in T which are not parametrically definable forms a subset of the set of emergent paths.

Note however that not all emergent paths need be undefinable. For example, the path A in the tree T from Example 30 is definable in T using the formula

$$\exists y_1 \exists y_2 \left(y_1 \neq y_2 \wedge \bigwedge_{i=1}^2 (x < y_i) \wedge \bigwedge_{i=1}^2 (\neg \exists z (x < z \wedge z < y_i)) \right).$$

7.3 On non-definable paths

First, we will show that if a path A in a tree T is not parametrically definable, then there exists a ‘ladder’ of paths in T , different from A , such that A cannot be distinguished from the path corresponding to the n -th step of the ‘ladder’ by using first-order formulae of quantifier rank n .

Lemma 31 *Let T be a tree and let A be a path that is not parametrically definable in T . For every $a \in A$ and for every $n \in \mathbb{N}$, there exists $b \in A$ and $c \in T \setminus A$ with $b, c \geq a$ and such that $b_{\leq} \equiv_n c_{\leq}$.*

Proof Let $a \in A$ and let $n \in \mathbb{N}$ but suppose to the contrary that $x_{\leq} \not\equiv_n y_{\leq}$ for every $x \in A$ and for every $y \in T \setminus A$ with $x, y \geq a$. Let τ_1, \dots, τ_m be all n -characteristics of empty tuples over the language of ordered sets. Let $U = \{i : x_{\leq} \models \tau_i \text{ for some } x \in A \text{ with } x \geq a\}$. Then for every x satisfying $x \geq a$, we have that $x_{\leq} \models \tau_i$ for some $i \in U$ if and only if $x \in A$. Let $\theta(u, y) = y \leq u$. Then A can be defined in $(T; a)$ using the formula

$$\varphi(x, a) := x < a \vee x \geq a \wedge \left(\bigvee_{i \in U} \tau_i^{\theta(u, a)} \right),$$

a contradiction. ⊥

Proposition 32 *Let α be a linear order containing a greatest element and suppose the first-order theory of α can be axiomatized using the sentence σ . Let T be a definably (uniformly) α -like tree containing only finitely many paths which are not parametrically definable. Then for every $n \in \mathbb{N}$, there exists a pathwise uniformly α -like tree S such that $S \equiv_n T$.*

Proof It suffices to prove the result for large n , so let $n \geq \text{qr}(\sigma) + 1$. Let A_1, \dots, A_k be the paths in T that are not parametrically definable and for every i , let $a_i \in A_i$ be such that $a_i \notin A_j$ for all j ($j \neq i$). By Lemma 31, for every i , there exists $b_i \in A_i$ and $c_i \in T \setminus A_i$ with $b_i, c_i \geq a_i$ and such that $(b_i)_{\leq} \equiv_n (c_i)_{\leq}$.

Let S be the tree obtained by taking the tree T and, for every i , replacing the subtree $(b_i)_{\leq}$ with the tree $S_i := (c_i)_{\leq}$. From the way S is constructed, every path in S will contain a leaf node and hence every path in S is definable using that leaf as parameter. Let $\theta(x, y) := x \leq y$ and define $\tau := \forall y (\text{leaf}(y) \rightarrow \sigma^{\theta(x, y)}(y))$. Note that $\text{qr}(\tau) = \text{qr}(\sigma) + 1$. By Proposition 11 we get $T \equiv_n S$ and since T satisfies τ then S also satisfies τ . Since every path in S contains a leaf then it follows that every path in S satisfies σ and hence S is pathwise uniformly α -like. \dashv

8 Concluding remarks

In this paper we have started a systematic study of the classes and first-order theories of trees, naturally associated, in terms of the paths occurring in those trees and the first-order theories of those paths, with a given class \mathcal{C} of linear orders. We have identified eight generally different \mathcal{C} -classes of trees and have classified their set-theoretic relationships and the respective relationships between their first-order theories in the general case, i.e., without any specific assumptions about the class \mathcal{C} . We have then obtained some general results about the axiomatizations of the first-order theories of some \mathcal{C} -classes of trees and have indicated the problems obstructing such general results for the other classes. These problems arise from the possible emergence of non-definable paths in the trees, that need not satisfy the first-order theory of \mathcal{C} , so we have started analyzing definable and undefinable paths in trees. In a follow-up paper we will apply these, and further results, to establish complete axiomatizations of the first-order theories of the classes of \mathcal{C} -trees for some important classes \mathcal{C} of linear orders.

9 Funding

This work was supported by the Department of Labour of South Africa and the National Research Foundation of South Africa.

References

- [1] Backofen, R., Rogers, J., Vijay-Shankar, K.: A First-Order Axiomatization of the Theory of Finite Trees. *Journal of Logic, Language and Information* (4), 5–39 (1995).
- [2] Doets, K.: Completeness and Definability: Applications of the Ehrenfeucht Game in Second-Order and Intensional Logic. Ph.D. thesis, University of Amsterdam (1987).
- [3] Ebbinghaus H.-D., J. Flum and W. Thomas, *Mathematical Logic*, Springer, 1996.
- [4] Goranko, V.: Trees and Finite Branching. Proceedings of the 2nd Panhellenic Logic Symposium, 91–101 (1999).
- [5] Goranko, V., Zanardo, A.: From linear to branching-time temporal logics: transfer of semantics and definability. *Logic Journal of the IGPL* (15), 53–76 (2007).
- [6] Rosenstein J., *Linear Orderings*, Academic Press, 1982.
- [7] Schmerl, J.: On \aleph_0 -categoricity and the theory of trees. *Fundamenta Mathematicae* (94), 121–128 (1977).