

**A STUDY ON THE EFFICACY OF A HOMOEOPATHIC
SIMILLIMUM REMEDY IN THE TREATMENT OF CHRONIC
ASTHMA IN ADULTS**

A dissertation submitted to the Faculty of Health Sciences,
Technikon Witwatersrand, Johannesburg, in partial fulfillment for the
degree of Master of Technology: Homoeopathy

by

Arnike Redelinghuys
(Student number: 9901688)

Supervisor: _____
Dr. S. C. van Es Date

Co-supervisor: _____
Dr. H. Niehaus Date

Research promotor _____
Dr. R. K. Razlog Date

DECLARATION

I, Arnike Redelinghuys, declare that this dissertation is original. It is being submitted to the Technikon Witwatersrand for the degree: Master of Technology: Homoeopathy. It has not been submitted previously to this or any other institution for the purpose of obtaining a qualification.

(Signature of Candidate)

_____ day of _____, 2004

To my Grandmother
Gré de Chavonnes Vrugt

ACKNOWLEDGEMENTS

Dr Sonya van Es

for supervision, homoeopathic input and greatly appreciated advice in this research

Dr Henriette Niehaus

for encouragement, enthusiasm and input as co-supervisor

Dr Radmila Razlog

for guidance, expertise, advice and support as research promotor

Dr Kanellie Hatzikonstandinou

for valuable assistance and advice during the first part of this research

Dr Robbie Potenza

for guidance and support in this research

Mr Leon Brits

for advice and willingness to make his consultation room available

Mrs Riëtte Eiselen and Mr Eugene Redelinghuys

for advice on computation and analysis of the research statistics

Mr Jan Pieter Bosman and Mrs Marieke Redelinghuys

for advice, editing and unfailing patience in proofreading

and

to family and friends, whose support over the past five years

has made the study of Homoeopathy possible

ABSTRACT

Chronic asthma is characterised by widespread inflammation and reversible narrowing of the bronchial airways. Inflammation of the bronchial mucous membrane renders it hyper-responsive to a variety of stimuli, resulting in wheezing, coughing, chest tightness and breathing difficulty. Asthma was first recognised by the Chinese, four thousand years ago and Hippocrates described it in the fourth century BC. Despite the amount of literature and information available on asthma, this disease is still on the increase world-wide.

This research study was undertaken to establish the efficacy of homoeopathic simillimum remedies in the reduction of the frequency and severity of asthmatic symptoms, in adult individuals who suffered from chronic asthma.

The research was carried out at the TWR health clinic at Doornfontein, from 1 June 2003 to 31 December 2003. Ten participants, eight males and two females, of all ethnic groups and between the ages of twenty-one and thirty-one, took part in this research. Each participant was interviewed during a homoeopathic consultation. Simillimum remedies were given and the participants were monitored over a period of fourteen weeks, at fortnightly follow-up consultations.

Each participant received his/her own peak flow meter and a peak flow rate chart (Appendix E). Morning and evening PEFr were obtained by using the peak flow meter. These readings were recorded, along with any relevant symptoms experienced and/or bronchodilating medication used that day. Participants were required to record the above-mentioned data for two weeks before the homoeopathic treatment commenced and for the remaining twelve weeks of the study.

The following results were obtained:

- Most participants reported less frequent and less severe acute asthmatic attacks, as well as fewer nocturnal asthmatic symptoms.
- Most participants reported an increased ability to participate in exercise without the need for bronchodilating medication during these activities.

- No statistically significant changes were found on the morning or evening peak expiratory flow rates recorded over fourteen weeks.
- No statistically significant change was found on the frequency of bronchodilating medication used over fourteen weeks.

The following conclusions may be drawn from the recorded results:

- Homoeopathic simillimum remedies appear to have been effective in the treatment of chronic asthma in adults.
- Homoeopathic simillimum remedies appear to have been effective in the reduction of acute asthmatic attacks and nocturnal asthmatic symptoms.
- Homoeopathic simillimum remedies appear to have been effective in increasing the participants' abilities to partake in physical exercise, as well as in decreasing their need for bronchodilating medicine during exercise.

TABLE OF CONTENTS

	Page
DECLARATION	ii
DEDICATION	iii
ACKNOWLEDGEMENTS	iv
ABSTRACT	v
TABLE OF CONTENTS	vii
LIST OF APPENDICES	xv
LIST OF FIGURES	xviii
LIST OF TABLES	xix

CHAPTER 1

INTRODUCTION

1.1	PROBLEM STATEMENT	1
1.2	HYPOTHESES	1
1.3	PURPOSE OF STUDY	2
1.4	IMPORTANCE OF THE PROBLEM	2
1.5	DELIMITATIONS	2
1.6	ASSUMPTIONS	3
1.7	CRITERIA GOVERNING THE ADMISSIBILITY OF THE DATA	3

CHAPTER 2

LITERATURE REVIEW

2.1	ASTHMA	4
2.1.1	Causes of asthma and aggravating factors	4
2.1.2	Extrinsic asthma	6
2.1.2.1	Atopic asthma	6
2.1.2.2	Exercise-induced asthma	6
2.1.2.3	Occupational asthma	6
2.1.3	Intrinsic asthma	7
2.1.3.1	Medicine induced asthma	7
2.1.3.2	Emotion or stress-induced asthma	8
2.2	ANATOMY OF A HEALTHY RESPIRATORY SYSTEM	8
2.2.1	Lung embryology	8
2.2.2	Structure of adult lung	9
2.2.2.1	Respiratory passages	9
2.2.2.2	Lungs	10
2.2.2.3	Mechanism of pulmonary ventilation	11
2.2.2.4	Respiratory centre	12
2.2.3	Histology of the respiratory tract	12
2.3	PATHOPHYSIOLOGY OF ASTHMA	13
2.3.1	Introduction	13
2.3.2	Macroscopic pathophysiology	13
2.3.3	Microscopic pathophysiology	14
2.3.3.1	Bronchial epithelial damage	14
2.3.3.2	Bronchial wall remodelling	14
2.3.4	Mucous production in asthma	15
2.3.4.1	Curshmann's spirals	15
2.3.4.2	Charcot-Leyden crystals	15
2.3.4.3	Creola bodies	15

2.3.5	Principles of the inflammatory response	16
2.3.5.1	Early and late phases of asthma and airway inflammation	16
2.3.5.2	Immune cells participating in inflammation	16
2.3.5.3	Inflammatory mediators	18
2.4	DIAGNOSING ASTHMA	20
2.4.1	Introduction	20
2.4.2	Signs and symptoms of asthma	20
2.4.2.1	Nocturnal asthma	20
2.4.3	Classification of asthma severity	21
2.4.4	Pulmonary function tests	22
2.4.4.1	Spirometry	22
2.4.4.2	Peak expiratory flow rate (PEFR)	22
2.4.5	Differential diagnosis	23
2.4.5.1	Chronic obstructive pulmonary disease (COPD)	23
2.4.5.2	Pulmonary eosinophilia	23
2.4.5.3	Viral respiratory tract infection	24
2.4.5.4	Other differential diagnosis	24
2.5	TREATMENT AND MANAGEMENT OF ASTHMA	25
2.5.1	Introduction	25
2.5.2	Gaining control and stepwise approach to management	25
2.5.3	Asthma education	27
2.5.4	Asthma action plan (Appendix F)	27
2.5.5	Non-drug management	28
2.5.6	Conventional drug treatment	29
2.5.6.1	Introduction	29
2.5.6.2	Bronchodilators	29
2.5.6.3	Corticosteroids (Prednisone and Methylprednisolone)	29
2.5.6.4	Theophylline (Uni-dur and Uniphyll)	30
2.5.6.5	Leukotriene receptor antagonists (Singulair)	30
2.5.7	Delivery devices for inhaled medication	30

2.6	HOMOEOPATHY	32
2.6.1	Origins and principles of homoeopathy	32
2.6.2	Constitution and susceptibility	33
2.6.3	Miasms	33
2.6.4	Homoeopathic Materia Medica	34
2.6.4.1	Provings	34
2.6.4.2	Potentization	35
2.6.4.3	Side effects of homoeopathic remedies	35
2.6.5	Homoeopathic case-taking and prescribing	36
2.6.5.1	Initial consultation	36
2.6.5.2	Follow-up consultation	37
2.6.5.3	Homoeopathic prescribing	37
2.6.6	Homoeopathic treatment of chronic asthma in adults	38
2.6.6.1	Introduction	38
2.6.6.2	Homoeopathic remedies for the treatment of chronic asthma in adults	39
2.6.7	Other research on the homoeopathic treatment of chronic asthma in adults	40

CHAPTER 3

MATERIALS AND METHODOLOGY

3.1	LIST OF MATERIALS AND EQUIPMENT	42
3.2	METHODOLOGY	43
3.2.1	Study design	43
3.2.2	Recruitment of participants	43
3.2.3	Investigative procedures	44
3.2.4	Homoeopathic medication and treatment protocol	45
3.3	DATA ANALYSIS	45

CHAPTER 4

RESULTS

4.1	INTRODUCTION	46
4.2	DESCRIPTIVE ANALYSIS	46
4.2.1	Peak expiratory flow rates and conventional medicine used	46
4.2.2	Weather data	50
4.3	ANALYTICAL RESULTS	50

CHAPTER 5

DISCUSSION OF RESULTS

5.1	CASE ONE	52
5.1.1	First consultation: 9 June 2003	52
5.1.2	First follow-up consultation: 14 July 2003	55
5.1.3	Second follow-up consultation: 28 July 2003	55
5.1.4	Third follow-up consultation: 11 August 2003	56
5.1.5	Fourth follow-up consultation: 25 August 2003	57
5.1.6	Fifth follow-up consultation: 8 September 2003	58
5.1.7	Sixth follow-up consultation: 22 September 2003	59
5.1.8	Overview of case one	60
5.2	CASE TWO	60
5.2.1	First consultation: 9 June 2003	60
5.2.2	First follow-up consultation: 14 July 2003	63
5.2.3	Second follow-up consultation: 28 July 2003	63
5.2.4	Third follow-up consultation: 11 August 2003	64
5.2.5	Fourth follow-up consultation: 25 August 2003	65

5.2.6	Fifth follow-up consultation: 8 September 2003	66
5.2.7	Sixth follow-up consultation: 22 September 2003	67
5.2.8	Overview of case two	68
5.3	CASE THREE	69
5.3.1	First consultation: 2 June 2003	69
5.3.2	First follow-up consultation: 14 July 2003	71
5.3.3	Second follow-up consultation: 28 July 2003	72
5.3.4	Third follow-up consultation: 18 August 2003	73
5.3.5	Fourth follow-up consultation: 1 September 2003	74
5.3.6	Fifth follow-up consultation: 15 September 2003	75
5.3.7	Sixth follow-up consultation: 29 September 2003	76
5.3.8	Overview of case three	77
5.4	CASE FOUR	77
5.4.1	First consultation: 25 June 2003	77
5.4.2	First follow-up consultation: 28 July 2003	80
5.4.3	Second follow-up consultation: 11 August 2003	81
5.4.4	Third follow-up consultation: 1 September 2003	82
5.4.5	Fourth follow-up consultation: 29 September 2003	83
5.4.6	Fifth follow-up consultation: 15 October 2003	84
5.4.7	Sixth follow-up consultation: 27 October 2003	86
5.4.8	Overview of case four	87
5.5	CASE FIVE	87
5.5.1	First consultation: 21 July 2003	87
5.5.2	First follow-up consultation: 18 August 2003	89
5.5.3	Second follow-up consultation: 1 September 2003	90
5.5.4	Third follow-up consultation: 15 September 2003	91
5.5.5	Fourth follow-up consultation: 29 September 2003	91
5.5.6	Fifth follow-up consultation: 13 October 2003	92
5.5.7	Sixth follow-up consultation: 27 October 2003	93

5.5.8	Overview of case five	94
5.6	CASE SIX	94
5.6.1	First consultation: 6 August 2003	94
5.6.2	First follow-up consultation: 8 September 2003	97
5.6.3	Second follow-up consultation: 22 September 2003	99
5.6.4	Third follow-up consultation: 8 October 2003	100
5.6.5	Fourth follow-up consultation: 20 October 2003	101
5.6.6	Fifth follow-up consultation: 5 November 2003	102
5.6.7	Sixth follow-up consultation: 17 November 2003	103
5.6.8	Overview of case six	104
5.7	CASE SEVEN	104
5.7.1	First consultation: 3 September 2003	104
5.7.2	First follow-up consultation: 29 September 2003	107
5.7.3	Second follow-up consultation: 13 October 2003	108
5.7.4	Third follow-up consultation: 5 November 2003	108
5.7.5	Fourth follow-up consultation: 19 November 2003	109
5.7.6	Fifth follow-up consultation: 1 December 2003	110
5.7.8	Overview of case seven	111
5.8	CASE EIGHT	111
5.8.1	First consultation: 21 August 2003	111
5.8.2	First follow-up consultation: 16 September 2003	114
5.8.3	Second follow-up consultation: 8 October 2003	115
5.8.4	Third follow-up consultation: 20 October 2003	116
5.8.5	Fourth follow-up consultation: 5 November 2003	117
5.8.6	Fifth follow-up consultation: 26 November 2003	118
5.8.7	Sixth follow-up consultation: 8 December 2003	119
5.8.8	Overview of case eight	120

5.9	CASE NINE	121
5.9.1	First consultation: 10 September 2003	121
5.9.2	First follow-up consultation: 7 October 2003	123
5.9.3	Second follow-up consultation: 21 October 2003	124
5.9.4	Third follow-up consultation: 5 November 2003	125
5.9.5	Fourth follow-up consultation: 19 November 2003	126
5.9.6	Fifth follow-up consultation: 1 December 2003	127
5.9.7	Overview of case nine	127
5.10	CASE TEN	128
5.10.1	First consultation: 22 September 2003	128
5.10.2	First follow-up consultation: 20 October 2003	130
5.10.3	Second follow-up consultation: 5 November 2003	131
5.10.4	Third follow-up consultation: 27 November 2003	132
5.10.5	Fourth follow-up consultation: 15 December 2003	133
5.10.6	Overview of case	133
CHAPTER 6		
CONCLUSIONS AND RECOMMENDATIONS		
6.1	CONCLUSION	135
6.2	RECOMMENDATIONS	136
6.2.1	Continued studies	136
6.2.2	Benefits of this study	137
6.2.3	Limitations of this study	137
REFERENCES		138

LIST OF APPENDICES

	Page
APPENDIX A:	
Patient consent and information form	147
APPENDIX B:	
Screening questionnaire	149
APPENDIX C:	
Questionnaire	150
APPENDIX D:	
Follow-up questionnaire	151
APPENDIX E:	
Peak expiratory flow rate chart	152
APPENDIX F:	
Asthma action plan	154
APPENDIX G:	
Electronic repertorisation sheet on common asthma symptoms (Witco and Stevenson, 1998)	155
APPENDIX H:	
Weather data	156
APPENDIX I:	
GLM repeated measures on morning PEFr test of within-subject contrast	161

APPENDIX J:	
GLM repeated measures on evening PEFr test of within subject contrast	162
APPENDIX K:	
GLM repeated measures on bronchodilating medication used in two weeks, test of within-subject contrast	163
APPENDIX L:	
Repertorisation for case one	164
APPENDIX M:	
Repertorisation for case two	165
APPENDIX N:	
Repertorisation for case three	166
APPENDIX O:	
Repertorisation for case four	167
APPENDIX P:	
Repertorisation for case five	168
APPENDIX Q:	
Repertorisation for case six	169
APPENDIX R:	
Repertorisation for case seven	170
APPENDIX S:	
Repertorisation for case eight	171

APPENDIX T:

Repertorisation for case nine 172

APPENDIX U:

Repertorisation for case ten 173

LIST OF FIGURES

	Page
Figure 4.2.1.1: Boxplot for morning PEFr	47
Figure 4.2.1.2: Boxplot for evening PEFr	48
Figure 4.2.1.3: Boxplot for bronchodilating medicine used in two weeks	49

LIST OF TABLES

	Page
Table 4.2.1.1: Descriptive statistics of morning PEFR (L/min)	47
Table 4.2.1.2: Descriptive statistics of evening PEFR (L/min)	48
Table 4.2.1.3: Descriptive statistics of bronchodilating medication used in two weeks	49
Table 4.3.1: Tests of within-subjects effects for PEFR	50